

PRODUCT MONOGRAPH

 EMEND[®] IV

fosaprepitant for injection

150 mg fosaprepitant /vial
(as fosaprepitant dimeglumine)

Neurokinin 1 (NK₁) receptor antagonist

Merck Canada Inc.
16750 route Transcanadienne
Kirkland, QC Canada H9H 4M7

Date of Revision:
January 30, 2012

www.merck.ca

Submission Control No: 150287

Table of Contents

PART I: HEALTH PROFESSIONAL INFORMATION.....3

SUMMARY PRODUCT INFORMATION3

INDICATIONS AND CLINICAL USE.....3

CONTRAINDICATIONS3

WARNINGS AND PRECAUTIONS.....4

ADVERSE REACTIONS.....5

DRUG INTERACTIONS9

DOSAGE AND ADMINISTRATION14

OVERDOSAGE15

ACTION AND CLINICAL PHARMACOLOGY16

STORAGE AND STABILITY19

DOSAGE FORMS, COMPOSITION AND PACKAGING19

PART II: SCIENTIFIC INFORMATION20

PHARMACEUTICAL INFORMATION.....20

CLINICAL TRIALS.....21

DETAILED PHARMACOLOGY26

TOXICOLOGY29

REFERENCES32

PART III: CONSUMER INFORMATION.....34

EMEND® IV

fosaprepitant for injection

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form / Strength	Clinically Relevant Non-Medicinal Ingredients
Intravenous	Lyophilized powder/ 150 mg/vial	<i>For a complete listing see DOSAGE FORMS, COMPOSITION AND PACKAGING section.</i>

INDICATIONS AND CLINICAL USE

EMEND® IV (fosaprepitant dimeglumine), in combination with a 5-HT₃ antagonist class of antiemetics and dexamethasone, is indicated for the prevention of acute and delayed nausea and vomiting due to highly emetogenic cancer chemotherapy.

Geriatrics (≥65 years of age): In clinical studies, the efficacy and safety of aprepitant in the elderly (≥65 years) were comparable to those seen in younger patients (<65 years). No dosage adjustment is necessary in elderly patients.

Pediatrics (<18 years of age): No data available.

CONTRAINDICATIONS

- Patients who are hypersensitive to EMEND® IV, aprepitant, polysorbate 80, or to any ingredient in the formulation. For a complete listing, see the DOSAGE FORMS, COMPOSITION AND PACKAGING section of the product monograph.
- EMEND® IV should not be used concurrently with pimozone, terfenadine, astemizole, or cisapride. Inhibition of cytochrome P450 isoenzyme 3A4 (CYP3A4) by aprepitant could result in elevated plasma concentrations of these drugs, potentially causing serious or life-threatening reactions (see DRUG INTERACTIONS).

WARNINGS AND PRECAUTIONS

Serious Warnings and Precautions

Drug interactions with:

- Medicinal products that are metabolized through CYP3A4 (see DRUG INTERACTIONS)
- Warfarin (see DRUG INTERACTIONS)
- Hormonal contraception (see DRUG INTERACTIONS)

The following are clinically significant adverse events:

- Immediate hypersensitivity reactions (see Hypersensitivity Reactions)

Hypersensitivity Reactions

Isolated reports of immediate hypersensitivity reactions including flushing, erythema, and dyspnea have occurred during infusion of fosaprepitant. These hypersensitivity reactions have generally responded to discontinuation of the infusion and administration of appropriate therapy. The infusion should not be reinitiated in patients who experience hypersensitivity reactions (see CONTRAINDICATIONS).

Special Populations

Pregnant Women: There are no adequate and well-controlled studies in pregnant women; therefore, EMEND[®] IV are not recommended for use during pregnancy unless clearly necessary (see TOXICOLOGY, Reproduction and Development).

Nursing Women: EMEND[®] IV, when administered intravenously, is rapidly converted to aprepitant. Aprepitant is excreted in the milk of lactating rats. It is not known whether this drug is excreted in human milk; therefore, breastfeeding is not recommended during treatment with EMEND[®] IV.

Pediatrics (<18 years of age): Safety and effectiveness of EMEND[®] IV in pediatric patients have not been established.

Geriatrics (≥65 years of age): In 2 well-controlled clinical studies, of the total number of patients (N=544) treated with aprepitant, 31% were 65 and over, while 5% were 75 and over. No overall differences in safety or effectiveness were observed between these subjects and younger subjects. Greater sensitivity of some older individuals cannot be ruled out. Dosage adjustment in the elderly is not necessary.

Hepatic Impairment: There are no clinical data or pharmacokinetic data in patients with severe hepatic impairment (Child-Pugh score >9). EMEND[®] IV should be used with caution in these patients (see ACTION AND CLINICAL PHARMACOLOGY, Special Populations and Conditions).

ADVERSE REACTIONS

Clinical Trial Adverse Experiences

Since EMEND[®] IV is converted to aprepitant, those adverse reactions associated with aprepitant are to be expected to occur with EMEND[®] IV. The overall safety of fosaprepitant was evaluated in approximately 1100 individuals and the overall safety of aprepitant was evaluated in approximately 6500 individuals. Injection site adverse reaction can also be expected with fosaprepitant.

Oral Aprepitant

Highly Emetogenic Chemotherapy (HEC)

In 2 well-controlled clinical trials in patients receiving cisplatin-based chemotherapy, 544 patients were treated with aprepitant during Cycle 1 of chemotherapy and 413 of these patients continued into the Multiple-Cycle extension for up to 6 cycles of chemotherapy. Oral aprepitant was given in combination with ondansetron and dexamethasone and was generally well tolerated. Most adverse experiences reported in these clinical studies were described as mild to moderate in intensity.

In Cycle 1, clinical adverse experiences were reported in approximately 74% of patients treated with the aprepitant regimen compared with approximately 72% of patients treated with standard therapy.

Table 1 shows the percent of patients with clinical adverse experiences reported at an incidence $\geq 3\%$.

Table 1 – All adverse experiences, regardless of causality, (incidence $\geq 3\%$) occurring in patients receiving highly emetogenic chemotherapy who were treated with the aprepitant regimen for chemotherapy induced nausea and vomiting (CINV) in clinical studies (cycle 1)

	Aprepitant Regimen N=544 %	Standard Therapy N=550 %
General Disorders and Administration Site Conditions		
Asthenia	10.1	7.1
Fatigue	8.8	5.8
Pyrexia	2.4	3.1
Gastrointestinal Disorders		
Abdominal Pain	4.2	3.1
Abdominal Pain Upper	3.7	3.3
Constipation	10.8	12.7
Diarrhea	10.3	7.5
Dyspepsia	8.3	6.9
Gastritis	3.9	2.9

	Aprepitant Regimen N=544 %	Standard Therapy N=550 %
Nausea	12.7	11.8
Stomatitis	2.2	3.1
Vomiting	7.5	7.5
Ear and Labyrinth Disorders		
Tinnitus	3.7	3.6
Metabolism and Nutrition Disorders		
Decreased appetite	11.8	10.0
Dehydration	5.5	4.7
Nervous System Disorders		
Dizziness	6.3	4.4
Headache	8.3	8.4
Psychiatric Disorders		
Insomnia	2.9	3.1
Respiratory Thoracic and Mediastinal Disorders		
Hiccups	10.8	5.6

In addition, isolated cases of serious adverse experiences, regardless of causality, of bradycardia, disorientation, and perforating duodenal ulcer were reported in highly emetogenic CINV clinical studies.

In an additional active-controlled clinical study (study P017L) in 1169 patients receiving aprepitant and highly emetogenic chemotherapy, the adverse experience profile was generally similar to that seen in the other HEC studies with aprepitant.

Additional Clinical Trial Adverse Experiences (less than 3% and considered to be clinically important), Regardless of Causality, Occurring in Patients Receiving aprepitant for CINV:

Blood and lymphatic system disorders: anemia, febrile neutropenia, neutropenia, thrombocytopenia.

Cardiac disorders: myocardial infarction, palpitations, tachycardia.

Eye disorders: conjunctivitis.

Gastrointestinal disorders: dry mouth, dysphagia, epigastric discomfort, eructation, flatulence, gastroesophageal reflux disease, odynophagia, salivary hypersecretion.

General disorders and administrative site conditions: chest pain, edema peripheral, malaise, pain.

Infections and infestations: oral candidiasis, pharyngitis, septic shock.

Investigations: weight decreased.

Metabolism and nutrition disorders: diabetes mellitus, hypokalemia.

Musculoskeletal and connective tissue disorders: musculoskeletal pain.

Nervous system disorders: dysgeusia, peripheral neuropathy, peripheral sensory neuropathy.

Psychiatric disorders: abnormal dreams, anxiety, confusion, depression, disorientation.

Renal and urinary disorders: dysuria, renal insufficiency.

Respiratory, thoracic and mediastinal disorders: cough, dyspnea, oropharyngeal pain, pneumonitis, pulmonary embolism, respiratory insufficiency, vocal disturbance.

Skin and subcutaneous tissue disorders: erythema, hyperhidrosis, acne, alopecia, rash.

Vascular disorders: deep venous thrombosis, flushing, hot flush, hypertension, hypotension.

Other Clinical Trials

Stevens-Johnson syndrome was reported as a serious adverse experience in a patient receiving aprepitant with cancer chemotherapy in another CINV study.

Abnormal Hematologic and Clinical Chemistry Findings

Table 2 shows the percent of patients with laboratory adverse experiences reported at an incidence $\geq 3\%$ in patients receiving highly emetogenic chemotherapy.

Table 2 – All laboratory abnormalities, regardless of causality, (incidence $\geq 3\%$) occurring in patients receiving highly emetogenic chemotherapy who were treated with the aprepitant regimen for CINV in clinical studies (cycle 1)

	Aprepitant Regimen	Standard Therapy
	N=544 %	N=550 %
ALT increased	5.9	3.8
Blood urea increased	4.6	3.5
Blood creatinine increased	3.7	4.3
Protein urine present	6.1	4.5

In other studies, laboratory adverse experiences reported at an incidence $\geq 3\%$ were decreased neutrophil counts (4.6% vs. 4.6%) and decreased white blood cell counts (5.1% vs. 4.7%) in patients receiving an aprepitant regimen or standard therapy, respectively.

Other Abnormal Hematological and Clinical Chemistry Findings Observed in Clinical Trials

The following additional laboratory adverse experiences, regardless of causality, were reported in patients treated with aprepitant regimen: AST increased, blood alkaline phosphatase increased, blood glucose increased, blood sodium decreased, white blood cell count increased, red blood cell urine positive, white blood cell urine positive. The adverse experiences of increased AST and ALT were generally mild and transient.

The adverse experience profiles in the Multiple-Cycle extensions of CINV studies for up to 6 cycles of chemotherapy were generally similar to those observed in Cycle 1.

Fosaprepitant (150 mg intravenous formulation)

In an active-controlled clinical study in patients receiving highly emetogenic chemotherapy, safety was evaluated for 1143 patients receiving the 1-day regimen of EMEND[®] IV 150 mg compared to 1169 patients receiving the 3-day regimen of EMEND[®] (aprepitant). The safety profile was generally similar to that seen in prior HEC studies with aprepitant (see Table 1). However, infusion-site reactions occurred at a higher incidence in the fosaprepitant group (3.0%) compared to those in the aprepitant group (0.5%), and there were significantly more cases of infusion-associated thrombophlebitis and infusion-site pain in the fosaprepitant group compared to those in the aprepitant group (thrombophlebitis: 0.8% vs. 0.1%; infusion-site pain: 1.4% vs. 0.1%). Infusion-site reactions included infusion-site erythema, infusion-site pruritus, infusion-site pain, infusion-site induration, and infusion-site thrombophlebitis.

In addition, there were more patients with adverse events of hypertension in the fosaprepitant group (1.5%) than in the aprepitant group (0.6%), as well as more patients with potential hypersensitivity adverse events in the fosaprepitant group (3.7%) than in the aprepitant group (3.1%).

Table 3 shows the percent of patients with clinical adverse experiences reported with fosaprepitant at an incidence >3%, regardless of causality.

Table 3 – All adverse experiences, regardless of causality, (incidence ≥3%) occurring in patients receiving highly emetogenic chemotherapy who were treated with fosaprepitant for CINV in a clinical study

	Fosaprepitant Regimen N=1143 %	Aprepitant Regimen N=1169 %
Blood and lymphatic system disorders		
Neutropenia	3.9	3.3
Gastrointestinal disorders		
Abdominal pain	3.1	3.3
Abdominal pain upper	4.0	2.6
Constipation	10.6	9.6
Diarrhea	7.8	8.7
Dyspepsia	4.4	3.3
Nausea	5.9	6.9
Vomiting	6.6	5.6

	Fosaprepitant Regimen N=1143 %	Aprepitant Regimen N=1169 %
General disorders and administration site conditions		
Asthenia	8.6	11.6
Fatigue	4.6	4.9
Metabolism and nutrition disorders		
Anorexia	6.6	9.1
Nervous system disorders		
Dizziness	3.3	3.0
Headache	4.0	4.1
Respiratory, thoracic, and mediastinal disorders		
Hiccups	5.6	6.3

Clinically significant laboratory analysis during the follow-up time period (Day 6 to 29) indicated a higher incidence of serum alanine aminotransferase >5X ULN in patients treated with the fosaprepitant regimen (1.8%) compared to patients treated with the aprepitant regimen (0.5%).

Post-Market Adverse Drug Reactions

Regardless of causality with aprepitant the following adverse events have been reported rarely or very rarely and occur with multiple confounding factors: loss of consciousness, depressed level of consciousness, convulsion, somnolence, paresthesia, syndrome of inappropriate antidiuretic hormone, hallucination, pruritus, rash, urticaria, Stevens-Johnson syndrome/toxic epidermal necrolysis, and hypersensitivity reactions including anaphylactic reactions.

Immediate hypersensitivity reactions have been observed during the infusion of fosaprepitant which may include the following: flushing, erythema, dyspnea (see WARNINGS AND PRECAUTIONS).

DRUG INTERACTIONS

Following the infusion of fosaprepitant 150 mg, a higher aprepitant C_{max} (~ 2.6-fold) was observed compared to oral aprepitant (125 mg). A theoretical risk for increased adverse experiences due to a higher peak aprepitant exposure cannot be ruled out.

Serious Drug Interactions

- Fosaprepitant is rapidly converted to aprepitant, which is an inhibitor of CYP3A4. Fosaprepitant should be used with caution in patients receiving concomitant medicinal products that are primarily metabolized through CYP3A4 and CYP2C9, including chemotherapy agents. Moderate inhibition of CYP3A4 by aprepitant and weak inhibition of CYP3A4 by fosaprepitant 150 mg could result in elevated plasma concentrations of these concomitant medicinal products. Induction of CYP2C9 by aprepitant could result in decreased plasma concentrations of these concomitant medicinal products (see CONTRAINDICATIONS and DRUG INTERACTIONS).
- The effect of oral aprepitant on the pharmacokinetics of orally administered CYP3A4 substrates is greater than the effect of oral aprepitant on the pharmacokinetics of intravenously administered CYP3A4 substrates.
- Coadministration of oral aprepitant with warfarin results in decreased prothrombin time, reported as International Normalized Ratio (INR). In patients on chronic warfarin therapy, the prothrombin time (INR) should be closely monitored in the 2-week period, particularly at 7 to 10 days, following initiation of fosaprepitant with each chemotherapy cycle (see DRUG INTERACTIONS).
- The efficacy of hormonal contraceptives during and for 28 days after administration of fosaprepitant may be reduced. Alternative or back-up methods of contraception should be used during treatment with fosaprepitant and for 1 month following administration (see DRUG INTERACTIONS).

Overview

Drug interactions following administration of fosaprepitant are likely to occur with drugs that interact with oral aprepitant. The following information was derived from studies conducted with oral aprepitant and studies conducted with fosaprepitant coadministered with dexamethasone, midazolam or diltiazem.

Aprepitant is a substrate, a moderate inhibitor, and an inducer of CYP3A4 when administered as a 3-day antiemetic dosing regimen for CINV. Aprepitant is also an inducer of CYP2C9.

Fosaprepitant 150 mg, given as a single dose, is a weak inhibitor of CYP3A4, and does not induce CYP3A4.

Chronic continuous use of EMEND[®] IV is not recommended because it has not been studied and because the drug interaction profile may change during chronic dosing.

Effect of fosaprepitant/aprepitant on the pharmacokinetics of other agents

Aprepitant, as a moderate inhibitor of CYP3A4, and fosaprepitant 150 mg, as a weak inhibitor of CYP3A4, can increase plasma concentrations of orally coadministered medicinal products that are metabolized through CYP3A4 (see CONTRAINDICATIONS). Aprepitant may increase the plasma concentration of orally administered CYP3A4 substrates to a greater extent than if the substrate was administered intravenously.

Aprepitant has been shown to induce the metabolism of S(-) warfarin and tolbutamide, which are metabolized through CYP2C9. Coadministration of fosaprepitant or oral aprepitant with these drugs or other drugs that are known to be metabolized by CYP2C9, such as phenytoin, may result in lower plasma concentrations of these drugs.

Effect of other agents on the pharmacokinetics of aprepitant

Aprepitant is a substrate for CYP3A4; therefore, coadministration of fosaprepitant with drugs that inhibit CYP3A4 activity may result in increased plasma concentrations of aprepitant.

Consequently, concomitant administration of fosaprepitant with strong CYP3A4 inhibitors (e.g., ketoconazole, itraconazole, nefazodone, troleandomycin, clarithromycin, ritonavir, nelfinavir) should be approached cautiously. Moderate CYP3A4 inhibitors (e.g., diltiazem) resulted in a 2-fold increase in plasma concentrations of aprepitant; therefore, concomitant administration should also be approached with caution.

Aprepitant is a substrate for CYP3A4; therefore, coadministration of fosaprepitant with drugs that strongly induce CYP3A4 activity (e.g., rifampin, carbamazepine, phenytoin) may result in reduced plasma concentrations and decreased efficacy.

Drug-Drug Interactions

Table 4 – Established or potential drug-drug interactions

Proper name	Ref	Effect	Clinical comment
pimozide	T	↑ pimozide concentration	Potentially causing serious or life-threatening reactions.
terfenadine	T	↑ terfenadine concentration	Potentially causing serious or life-threatening reactions.
Astemizole	T	↑ astemizole concentration	Potentially causing serious or life-threatening reactions.
Cisapride	T	↑ cisapride concentration	Potentially causing serious or life-threatening reactions.
Warfarin	CT	↓ Warfarin concentration ↓ INR	In patients on chronic warfarin therapy, the INR should be closely monitored in the 2-week period, particularly at 7 to 10 days, following initiation of fosaprepitant with each chemotherapy cycle (see WARNINGS AND PRECAUTIONS and DETAILED PHARMACOLOGY).
tolbutamide	CT	↓ tolbutamide concentration	Aprepitant induces the metabolism of drug metabolized by CYP2C9 (see DETAILED PHARMACOLOGY).

Proper name	Ref	Effect	Clinical comment
Phenytoin	T	↓ phenytoin concentration	Aprepitant induces the metabolism of drug metabolized by CYP2C9.
dexamethasone	CT	↑ dexamethasone concentration	The usual oral dexamethasone dose on Days 1 and 2 should be reduced by approximately 50% when coadministered with fosaprepitant 150 mg IV on Day 1 to achieve exposures of dexamethasone similar to those obtained when it is given without fosaprepitant 150 mg (see DETAILED PHARMACOLOGY).
methylprednisolone	CT	↑ methylprednisolone concentration	See DETAILED PHARMACOLOGY.
hormone contraceptives with all routes of administration	CT	↓ hormone concentration	The efficacy of hormonal contraceptives during and for 28 days after administration fosaprepitant may be reduced. Alternative or back-up methods of contraception should be used during treatment with fosaprepitant and for 1 month following administration (see WARNINGS AND PRECAUTIONS and DETAILED PHARMACOLOGY).
Midazolam oral and IV	CT	↑ midazolam concentration	The potential effects of increased plasma concentrations of midazolam or other benzodiazepines metabolized via CYP3A4 (alprazolam, triazolam) should be considered when coadministering these agents with fosaprepitant. Fosaprepitant 150 mg IV as a single dose is a weak CYP3A4 inhibitor on Day 1 with no evidence of inhibition or induction of CYP3A4 observed on Day 4 (see DETAILED PHARMACOLOGY).
ketoconazole	CT	↑ aprepitant concentration	Concomitant administration of fosaprepitant with strong CYP3A4 inhibitors should be approached cautiously (see DETAILED PHARMACOLOGY).
Rifampin	CT	↓ aprepitant concentration	Coadministration of fosaprepitant with drugs that induce CYP3A4 activity may result in reduced plasma concentrations and decreased efficacy of EMEND® IV (see DETAILED PHARMACOLOGY).
Diltiazem	CT	↑ aprepitant and diltiazem concentration	No clinically meaningful changes in ECG, heart rate, PR interval or blood pressure beyond those changes induced by diltiazem alone (see DETAILED PHARMACOLOGY).
paroxetine	CT	↓ aprepitant and paroxetine concentration	See DETAILED PHARMACOLOGY.

Legend: CT = Clinical Trial; T = Theoretical

Fosaprepitant is unlikely to interact with drugs that are substrates for the P-glycoprotein transporter, as demonstrated by the lack of interaction of oral aprepitant with digoxin in a clinical drug interaction study.

5-HT₃ antagonists: In clinical drug interaction studies, aprepitant did not have clinically important effects on the pharmacokinetics of ondansetron administered intravenously, granisetron administered orally, or hydrodolasetron (the active metabolite of dolasetron) following oral administration of dolasetron.

Chemotherapeutic agents: Chemotherapy agents that are known to be metabolized by CYP3A4 include docetaxel, paclitaxel, etoposide, irinotecan, ifosfamide, imatinib, vinorelbine, vinblastine and vincristine. In clinical studies, the oral aprepitant regimen was administered with the following chemotherapeutic agents metabolized primarily or in part by CYP3A4: etoposide, vinorelbine, docetaxel, paclitaxel, irinotecan and ifosfamide. The doses of these agents were not adjusted to account for potential drug interactions. However, caution is advised and additional monitoring may be appropriate in patients receiving chemotherapy agents known to be metabolized by CYP3A4, especially those not studied in the clinical trials, including vinblastine, vincristine and imatinib (see WARNINGS AND PRECAUTIONS).

Docetaxel: In a separate pharmacokinetic study, oral aprepitant did not influence the pharmacokinetics of docetaxel.

Etoposide, paclitaxel: No pharmacokinetic studies to determine the effect of oral aprepitant on the concentration of etoposide or paclitaxel were performed.

Vinorelbine: In a separate pharmacokinetic study, oral aprepitant (125 mg/80 mg regimen) did not influence the pharmacokinetics of vinorelbine.

Drug-Food Interactions

EMEND[®] IV may be administered with or without food.

Drug-Herb Interactions

Interactions with herbal products have not been established.

Drug-Laboratory Interactions

Interactions with laboratory tests have not been established.

DOSAGE AND ADMINISTRATION

Dosing Consideration

EMEND[®] IV is available as a 150 mg for IV infusion.

EMEND[®] IV has not been demonstrated to be effective as a single anti-emetic agent and must be administered with other anti-emetic agents.

Recommended Dose and Dosage Adjustment

EMEND[®] IV 150 mg is administered on Day 1 only as an infusion **over 20–30 minutes** initiated approximately 30 minutes prior to chemotherapy. EMEND[®] IV should be administered in conjunction with a corticosteroid and a 5-HT₃ antagonist as specified in the table below.

Table 5 – Recommended dosing for the prevention of nausea and vomiting associated with highly emetogenic cancer chemotherapy

	Day 1	Day 2	Day 3	Day 4
EMEND [®] IV	150 mg IV	none	none	none
Dexamethasone*	12 mg orally	8 mg orally	8 mg orally bid	8 mg orally bid
Ondansetron [†]	32 mg IV	none	none	none

* Dexamethasone should be administered 30 minutes prior to chemotherapy treatment on Day 1 and in the morning on Days 2 through 4. Dexamethasone should also be administered in the evenings on Days 3 and 4. The dose of dexamethasone accounts for drug interactions.

[†] Ondansetron should be administered 30 minutes prior to chemotherapy treatment on Day 1.

For highly emetogenic chemotherapy, there is only limited efficacy data with EMEND[®] in combination with oral ondansetron or other 5-HT₃ antagonist class of antiemetics and dexamethasone.

Dosage Adjustment

See DRUG INTERACTIONS for additional information on the administration of fosaprepitant or aprepitant with corticosteroids.

Refer to each product's respective Product Monograph for additional information on coadministered antiemetic agents.

No dosage adjustment is necessary for the elderly.

No dosage adjustment is necessary based on gender or race.

No dosage adjustment is necessary for patients with severe renal insufficiency (creatinine clearance <30 mL/min) or for patients with end stage renal disease undergoing hemodialysis.

No dosage adjustment is necessary for patients with mild to moderate hepatic insufficiency (Child-Pugh score 5 to 9). There are no clinical data in patients with severe hepatic insufficiency (Child-Pugh score >9).

Administration

EMEND[®] IV is for IV infusion only, upon reconstitution and dilution and for single use only.

Instructions for reconstitution and dilution

1. Aseptically inject 5 mL 0.9% Sodium Chloride for injection (saline) into the vial. Assure that saline is added to the vial along the vial wall in order to prevent foaming. Swirl the vial gently. Avoid shaking and jetting saline into the vial.
2. Aseptically prepare an infusion bag filled with **145 mL** 0.9% NaCl for injection
3. Aseptically withdraw the entire volume from the vial and transfer it into an infusion bag containing 145 mL of saline to **yield a total volume of 150 mL** and a final concentration of approximately 1 mg fosaprepitant/mL. Gently invert the bag 2–3 times.

The reconstituted and diluted solutions should be used immediately; however, after reconstitution and dilution the final drug solution is stable for 24 hours at ambient room temperature (at or below 25°C).

Reconstituted and diluted solutions should be inspected for discoloration, cloudiness and particulate matter before administration whenever solution and container permit. Discard unused portion.

EMEND[®] IV is incompatible with any solutions containing divalent cations (e.g., Ca²⁺, Mg²⁺), including Hartman's and Lactated Ringer's Solution. EMEND[®] IV must not be reconstituted or mixed with solutions for which physical and chemical compatibility have not been established.

OVERDOSAGE

For management of a suspected drug overdose, contact your regional Poison Control Centre.

No specific information is available on the treatment of overdose. Single doses up to 200 mg of fosaprepitant IV and 600 mg of aprepitant were generally well tolerated in healthy subjects. Three out of 33 subjects receiving 200 mg of fosaprepitant experienced mild injection site thrombosis. Aprepitant was generally well tolerated when administered as 375 mg once daily for up to 42 days

to patients in non-CINV studies. In 33 cancer patients, administration of a single 375-mg dose of aprepitant on Day 1 and 250 mg once daily on Days 2 to 5 was generally well tolerated.

Drowsiness and headache were reported in one patient who ingested 1440 mg of aprepitant.

In the event of overdose, EMEND® IV should be discontinued and general supportive treatment and monitoring should be provided. Because of the antiemetic activity of aprepitant, drug-induced emesis may not be effective.

Aprepitant cannot be removed by hemodialysis.

ACTION AND CLINICAL PHARMACOLOGY

Mechanism of Action

Fosaprepitant is a prodrug of aprepitant and accordingly, its antiemetic effects are attributable to aprepitant.

Aprepitant has a unique mode of action; it is a selective high affinity antagonist at human substance P neurokinin 1 (NK₁) receptors. Counter-screening assays showed that aprepitant was at least 3,000-fold selective for the NK₁ receptor over other enzyme, transporter, ion channel and receptor sites including the dopamine and serotonin receptors that are targets for existing chemotherapy induced nausea and vomiting (CINV) therapies.

NK₁-receptor antagonists have been shown pre-clinically to inhibit emesis induced by cytotoxic chemotherapeutic agents, such as cisplatin, via central actions. Preclinical and human Positron Emission Tomography (PET) studies with aprepitant have shown that it is brain penetrant and occupies brain NK₁ receptors. Preclinical studies show that aprepitant has a long duration of central activity, inhibits both the acute and delayed phases of cisplatin-induced emesis, and augments the antiemetic activity of the 5-HT₃-receptor antagonist ondansetron and the corticosteroid dexamethasone against cisplatin-induced emesis.

Pharmacokinetics

Table 6 – Summary of pharmacokinetic parameters of EMEND® in healthy subjects

	C_{max} (µg/mL)	AUC_{0-24hr} (µg•hr/mL)
Day 1 oral dose aprepitant 125 mg	1.5	19.5
Day 3 oral dose aprepitant 80 mg	1.4	20.1

Absorption: Following a single intravenous 150-mg dose of fosaprepitant administered as a 20-minute infusion to healthy volunteers, the mean $AUC_{0-\infty}$ of aprepitant was 37.38 (\pm 14.75) mcg•hr/mL and the mean maximal aprepitant concentration was 4.15 (\pm 1.15) mcg/mL.

Fosaprepitant Administration

The mean absolute oral bioavailability of aprepitant is approximately 60 to 65% and the mean peak plasma concentration (C_{max}) of aprepitant occurred at approximately 4 hours (T_{max}). Oral administration of the capsule with a standard breakfast had no clinically meaningful effect on the bioavailability of aprepitant.

The pharmacokinetics of aprepitant are non-linear across the clinical dose range. In healthy young adults, the increase in $AUC_{0-\infty}$ was 26% greater than dose proportional between 80-mg and 125-mg single doses administered in the fed state.

Following oral administration of a single 125-mg dose of EMEND[®] on Day 1 and 80 mg once daily on Days 2 and 3, the AUC_{0-24hr} was approximately 19.5 μ g•hr/mL and 20.1 μ g•hr/mL on Day 1 and Day 3, respectively. The C_{max} of 1.5 μ g/mL and 1.4 μ g/mL were reached in approximately 4 hours (T_{max}) on Day 1 and Day 3, respectively.

Distribution: Fosaprepitant is rapidly converted to aprepitant.

Aprepitant is greater than 95% bound to plasma proteins. The geometric mean apparent volume of distribution at steady state (V_{dss}) is approximately 66 L in humans.

Aprepitant crosses the placenta in rats, and crosses the blood brain barrier in rats and ferrets. PET studies in humans indicate that aprepitant crosses the blood brain barrier (see ACTION AND CLINICAL PHARMACOLOGY).

Metabolism: Fosaprepitant was rapidly converted to aprepitant in *in vitro* incubations with liver preparations from nonclinical species (rat and dog) and humans. Furthermore, fosaprepitant underwent rapid and nearly complete conversion to aprepitant in S9 preparations from multiple other human tissues including kidney, lung and ileum. Thus, it appears that the conversion of fosaprepitant to aprepitant can occur in multiple extrahepatic tissues in addition to the liver. In humans, fosaprepitant administered intravenously was rapidly converted to aprepitant within 30 minutes following the end of infusion.

Aprepitant undergoes extensive metabolism. In healthy young adults, aprepitant accounts for approximately 24% of the radioactivity in plasma over 72 hours following a single oral 300-mg dose of [¹⁴C]-aprepitant, indicating a substantial presence of metabolites in the plasma. Seven metabolites of aprepitant, which are only weakly active, have been identified in human plasma. The metabolism of aprepitant occurs largely via oxidation at the morpholine ring and its side chains. *In vitro* studies using human liver microsomes indicate that aprepitant is metabolized primarily by CYP3A4 with minor metabolism by CYP1A2 and CYP2C19, and no metabolism by CYP2D6, CYP2C9, or CYP2E1.

All metabolites observed in urine, feces and plasma following an intravenous 100-mg [¹⁴C]-fosaprepitant dose were also observed following an oral dose of [¹⁴C]-aprepitant. Upon conversion of 245.3 mg of fosaprepitant dimeglumine (equivalent to 150 mg fosaprepitant free acid) to aprepitant, 23.9 mg of phosphoric acid and 95.3 mg of meglumine are liberated.

Excretion: Following administration of a single IV 100-mg dose of [¹⁴C]-fosaprepitant to healthy subjects, 57% of the radioactivity was recovered in urine and 45% in feces.

Aprepitant is eliminated primarily by metabolism; aprepitant is not renally excreted. Following administration of a single oral 300-mg dose of [¹⁴C]-aprepitant to healthy subjects, 5% of the radioactivity was recovered in urine and 86% in feces.

The apparent plasma clearance of aprepitant ranged from approximately 60 to 84 mL/min. The apparent terminal half-life ranged from approximately 9 to 13 hours.

Special Populations and Conditions

Pediatrics: The pharmacokinetics of EMEND[®] and EMEND[®] IV have not been evaluated in patients below 18 years of age.

Geriatrics: Following oral administration of a single 125-mg dose of EMEND[®] on Day 1 and 80 mg once daily on Days 2 through 5, the AUC_{0-24hr} of aprepitant was 21% higher on Day 1 and 36% higher on Day 5 in elderly (≥65 years) relative to younger adults. The C_{max} was 10% higher on Day 1 and 24% higher on Day 5 in elderly relative to younger adults. These differences are not considered clinically meaningful. No dosage adjustment for EMEND[®] is necessary in elderly patients.

Gender: Following oral administration of a single 125-mg dose of aprepitant, the C_{max} for aprepitant is 16% higher in females as compared with males. The half-life of aprepitant is 25% lower in females as compared with males and its T_{max} occurs at approximately the same time. No dosage adjustment is necessary based on gender.

Race: Following oral administration of a single 125-mg dose of aprepitant, the AUC_{0-24hr} is approximately 25% and 29% higher in Hispanics as compared with Caucasians and Blacks, respectively. The C_{max} is 22% and 31% higher in Hispanics as compared with Caucasians and Blacks, respectively. These differences are not considered clinically meaningful. No dosage adjustment is necessary based on race.

Hepatic Insufficiency: Fosaprepitant is metabolized in various extrahepatic tissues; therefore hepatic insufficiency is not expected to alter the conversion of fosaprepitant to aprepitant.

Oral aprepitant was well tolerated in patients with mild to moderate hepatic insufficiency. Following administration of a single 125-mg dose of oral aprepitant on Day 1 and 80 mg once daily on Days 2 and 3 to patients with mild hepatic insufficiency (Child-Pugh score 5 to 6), the AUC_{0-24hr} of aprepitant was 11% lower on Day 1 and 36% lower on Day 3, as compared with

healthy subjects given the same regimen. In patients with moderate hepatic insufficiency (Child-Pugh score 7 to 9), the AUC_{0-24hr} of Aprepitant was 10% higher on Day 1 and 18% higher on Day 3, as compared with healthy subjects given the same regimen. These differences in AUC_{0-24hr} are not considered clinically meaningful; therefore, no dosage adjustment is necessary in patients with mild to moderate hepatic insufficiency.

There are no clinical or pharmacokinetic data in patients with severe hepatic insufficiency (Child-Pugh score >9).

Renal Insufficiency: A single 240-mg dose of Aprepitant was administered to patients with severe renal insufficiency ($CrCl < 30$ mL/min) and to patients with end stage renal disease (ESRD) requiring hemodialysis.

In patients with severe renal insufficiency, the $AUC_{0-\infty}$ of total Aprepitant (unbound and protein bound) decreased by 21% and C_{max} decreased by 32%, relative to healthy subjects. In patients with ESRD undergoing hemodialysis, the $AUC_{0-\infty}$ of total Aprepitant decreased by 42% and C_{max} decreased by 32%. Due to modest decreases in protein binding of Aprepitant in patients with renal disease, the AUC of pharmacologically active unbound drug was not significantly affected in patients with renal insufficiency compared with healthy subjects. Hemodialysis conducted 4 or 48 hours after dosing had no significant effect on the pharmacokinetics of Aprepitant; less than 0.2% of the dose was recovered in the dialysate.

No dosage adjustment is necessary for patients with severe renal insufficiency or for patients with ESRD undergoing hemodialysis.

STORAGE AND STABILITY

Vials: store at 2–8°C.

DOSAGE FORMS, COMPOSITION AND PACKAGING

EMEND[®] IV is available as a 150 mg single dose per 10 mL glass vial as a white to off-white lyophilized solid to be administered intravenously as an infusion, upon reconstitution and dilution. One vial per carton.

Active ingredients:

Each vial of EMEND[®] IV 150 mg for intravenous administration contains 245.3 mg of fosaprepitant dimeglumine equivalent to 150 mg of fosaprepitant free acid.

Inactive ingredients:

Each vial of EMEND[®] IV for intravenous administration contains the following inactive ingredients: edetate disodium, polysorbate 80, lactose anhydrous, sodium hydroxide and/or hydrochloric acid (for pH adjustment).

CLINICAL TRIALS

Fosaprepitant, a prodrug of aprepitant, when administered intravenously is rapidly converted to aprepitant.

The pivotal efficacy studies were conducted with oral aprepitant. Oral administration of EMEND[®] (aprepitant) in combination with ondansetron and dexamethasone has been shown to prevent nausea and vomiting associated with highly emetogenic chemotherapy in well-controlled clinical studies. An active-controlled non-inferiority study comparing EMEND[®] IV 150 mg with oral aprepitant demonstrated efficacy of EMEND[®] IV 150 mg in the prevention of nausea and vomiting associated with highly emetogenic chemotherapy.

Study Demographics and Trial Design

Table 7 – Summary of patient demographics for clinical trials in highly emetogenic chemotherapy (HEC)

Study #	Trial design	Dosage, route of administration and duration	Study subjects (n=number)	Mean age (Range)	Gender
052 (HEC)	Randomized, double-blind, placebo-controlled, parallel-group	EMEND [®] 125 mg on Day 1 80 mg/day on Days 2 and 3 in combination with ondansetron 32 mg IV on Day 1 and dexamethasone 12 mg on Day 1 and 8 mg once daily on Days 2 through 4. OR Standard therapy which consisted of placebo in combination with ondansetron 32 mg IV on Day 1 and dexamethasone 20 mg on Day 1 and 8 mg twice daily on Days 2 through 4.	534	14–84	Male: 266 Female: 268
054 (HEC)	Randomized, double-blind, placebo-controlled, parallel-group	EMEND [®] 125 mg on Day 1 80 mg/day on Days 2 and 3 in combination with ondansetron 32 mg IV on Day 1 dexamethasone 12 mg on Day 1 and 8 mg once daily on Days 2 through 4. OR Standard therapy which consisted of placebo in combination with	569	18–82	Male: 283 Female: 286

Study #	Trial design	Dosage, route of administration and duration	Study subjects (n=number)	Mean age (Range)	Gender
		ondansetron 32 mg IV on Day 1 and dexamethasone 20 mg on Day 1 and 8 mg twice daily on Days 2 through 4.			
P017L (HEC)	Randomized, double-blind, active-controlled, parallel group, non-inferiority	Fosaprepitant 150 mg on Day 1 in combination with ondansetron 32 mg IV on Day 1 dexamethasone 12 mg on Day 1 and 8 mg on Day 2, and 8 mg twice daily on Days 3 and 4. OR Aprepitant 125 mg on Day 1 80 mg/day on Days 2 and 3 in combination with ondansetron 32 mg IV on Day 1 dexamethasone 12 mg on Day 1 and 8 mg once daily on Days 2 through 4.	2322	55 (19–86)	Male:1470 Female: 852

Oral administration of EMEND[®] (aprepitant) in combination with ondansetron and dexamethasone has been shown to prevent nausea and vomiting associated with highly emetogenic chemotherapy in well-controlled clinical studies.

3-Day Regimen of EMEND[®]

In the clinical studies 052 and 054 above, all enrolled patients received high-dose cisplatin ≥ 70 mg/m². Approximately 95% of the patients in the aprepitant group received a concomitant chemotherapeutic agent. The most common chemotherapeutic agents and the number of aprepitant patients exposed follows: etoposide (106), fluorouracil (100), gemcitabine (89), vinorelbine (82), paclitaxel (52), cyclophosphamide (50), doxorubicin (38), docetaxel (11). The efficacy of EMEND[®] has not been investigated in highly emetogenic chemotherapy clinical trials without cisplatin.

The antiemetic activity of EMEND[®] was evaluated during the acute phase (0 to 24 hours post-cisplatin treatment), the delayed phase (25 to 120 hours post-cisplatin treatment) and overall (0 to 120 hours post-cisplatin treatment) in Cycle 1. Efficacy was based on evaluation of the following endpoints:

Primary endpoint:

- complete response (defined as no emetic episodes and no use of rescue therapy)

Other prespecified endpoints:

- complete protection (defined as no emetic episodes, no use of rescue therapy, and a maximum nausea visual analogue scale [VAS] score <25 mm on a 0 to 100 mm scale)
- no emesis (defined as no emetic episodes regardless of use of rescue therapy)
- no nausea (maximum VAS <5 mm on a 0 to 100 mm scale)
- no significant nausea (maximum VAS <25 mm on a 0 to 100 mm scale)

A summary of the key study results from each individual study analysis is shown in Table 8 and in Table 9.

Table 8 – Percent of patients receiving highly emetogenic chemotherapy responding by treatment group and phase for study 1 – Cycle 1

ENDPOINTS	Aprepitant Regimen (N=260) [†] %	Standard Therapy (N=261) [†] %	p-Value
PRIMARY ENDPOINT			
Complete Response			
Overall [‡]	73	52	<0.001
OTHER PRESPECIFIED ENDPOINTS			
Complete Response			
Acute phase [§]	89	78	<0.001
Delayed phase	75	56	<0.001
Complete Protection			
Overall	63	49	0.001
Acute phase	85	75	NS*
Delayed phase	66	52	<0.001
No Emesis			
Overall	78	55	<0.001
Acute phase	90	79	0.001
Delayed phase	81	59	<0.001
No Nausea			
Overall	48	44	NS**
Delayed phase	51	48	NS**
No Significant Nausea			
Overall	73	66	NS**
Delayed phase	75	69	NS**

[†] N: Number of patients (older than 18 years of age) who received cisplatin, study drug, and had at least one post-treatment efficacy evaluation.

[‡] Overall: 0 to 120 hours post-cisplatin treatment.

[§] Acute phase: 0 to 24 hours post-cisplatin treatment.

^{||} Delayed phase: 25 to 120 hours post-cisplatin treatment.

* Not statistically significant when adjusted for multiple comparisons.

** Not statistically significant.

Visual analogue scale (VAS) score range: 0 mm = no nausea; 100 mm = nausea as bad as it could be.

Table 9 – Percent of patients receiving highly emetogenic chemotherapy responding by treatment group and phase for study 2 – Cycle 1

ENDPOINTS	Aprepitant Regimen (N=261) [†] %	Standard Therapy (N=263) [†] %	p-Value
PRIMARY ENDPOINT			
Complete Response			
Overall [‡]	63	43	<0.001
OTHER PRESPECIFIED ENDPOINTS			
Complete Response			
Acute phase [§]	83	68	<0.001
Delayed phase	68	47	<0.001
Complete Protection			
Overall	56	41	<0.001
Acute phase	80	65	<0.001
Delayed phase	61	44	<0.001
No Emesis			
Overall	66	44	<0.001
Acute phase	84	69	<0.001
Delayed phase	72	48	<0.001
No Nausea			
Overall	49	39	NS*
Delayed phase	53	40	NS*
No Significant Nausea			
Overall	71	64	NS**
Delayed phase	73	65	NS**

[†] N: Number of patients (older than 18 years of age) who received cisplatin, study drug, and had at least one post-treatment efficacy evaluation.

[‡] Overall: 0 to 120 hours post-cisplatin treatment.

[§] Acute phase: 0 to 24 hours post-cisplatin treatment.

^{||} Delayed phase: 25 to 120 hours post-cisplatin treatment.

* Not statistically significant when adjusted for multiple comparisons.


** Not statistically significant.

Visual analogue scale (VAS) score range: 0 mm = no nausea; 100 mm = nausea as bad as it could be.

In both studies, a statistically significantly higher proportion of patients receiving the aprepitant regimen in Cycle 1 had a complete response (primary endpoint), compared with patients receiving standard therapy. A statistically significant difference in complete response in favor of the aprepitant regimen was also observed when the acute phase and the delayed phase were analyzed separately.

In both studies, the estimated time to first emesis after initiation of cisplatin treatment was longer with the aprepitant regimen, and the incidence of first emesis was reduced in the aprepitant regimen group compared with standard therapy group as depicted in the Kaplan-Meier curves in Figure 1.

Figure 1 – Percent of patients receiving highly emetogenic chemotherapy who remain emesis free over time – Cycle 1



p-Value <0.001 based on a log rank test for Study 1 and Study 2; nominal p-values not adjusted for multiplicity.

Patient-Reported Outcomes: The impact of nausea and vomiting on patients' daily lives was assessed in Cycle 1 of both Phase III studies using the Functional Living Index–Emesis (FLIE), a validated nausea and vomiting-specific patient-reported outcome measure. Minimal or no impact of nausea and vomiting on patients' daily lives is defined as a FLIE total score >108. In each of the 2 studies, a higher proportion of patients receiving the aprepitant regimen reported minimal or no impact of nausea and vomiting on daily life (Study 1: 74% versus 64%; Study 2: 75% versus 64%).

Multiple-Cycle Extension: In the same 2 clinical studies, 851 patients continued into the Multiple-Cycle extension for up to 6 cycles of chemotherapy. The efficacy of the aprepitant regimen was maintained during all cycles.

1-Day Regimen of EMEND® IV

In a randomized, parallel, double-blind, active-controlled non-inferiority study, fosaprepitant 150 mg (N=1147) was compared with a 3-day aprepitant regimen (N=1175) in patients receiving a highly emetogenic chemotherapy regimen that included cisplatin (≥ 70 mg/m²). Other concomitant chemotherapy agents were administered similar to those in prior HEC studies described above. The fosaprepitant regimen consisted of fosaprepitant 150 mg on Day 1 in combination with ondansetron 32 mg IV on Day 1 and dexamethasone 12 mg on Day 1, 8 mg on Day 2, and 8 mg twice daily on Days 3 and 4. The aprepitant regimen consisted of aprepitant 125 mg on Day 1 and 80 mg/day on Days 2 and 3 in combination with ondansetron 32 mg IV on Day 1 and dexamethasone 12 mg on Day 1 and 8 mg daily on Days 2 through 4. Fosaprepitant placebo, aprepitant placebo, and dexamethasone placebo (in the evenings on Days 3 and 4) were used to maintain blinding.

Efficacy was based on the evaluation of the following composite measures: complete response in both the overall and delayed phases and no vomiting in the overall phase. The pre-specified non-inferiority margins were as follows: complete response in the overall phase -7 percentage points; complete response in the delayed phase -7.3 percentage points; and no vomiting in the overall phase -8.2 percentage points. EMEND[®] IV 150 mg was shown to be non-inferior to that of the 3-day regimen of aprepitant. A summary of the results is shown in Table 10.

Table 10 – Percent of Patients Receiving Highly Emetogenic Chemotherapy Responding by Treatment Group and Phase — Cycle 1			
ENDPOINTS*	Fosaprepitant Regimen (N=1106)** %	Aprepitant Regimen (N=1134)** %	Difference [†] (95% CI)
Complete Response[‡]			
Overall[§]	71.9	72.3	-0.4 (-4.1, 3.3)
Delayed phase ^{§§}	74.3	74.2	0.1 (-3.5, 3.7)
No Vomiting			
Overall [§]	72.9	74.6	-1.7 (-5.3, 2.0)

* Primary endpoint is bolded.

** N: Number of patients included in the primary analysis of complete response.

† Difference and confidence interval (CI) were calculated using the method proposed by Miettinen and Nurminen and adjusted for Gender.

‡ Complete Response: no vomiting and no use of rescue therapy.

§ Overall: 0 to 120 hours post-initiation of cisplatin chemotherapy.

§§ Delayed phase: 25 to 120 hours post-initiation of cisplatin chemotherapy.

DETAILED PHARMACOLOGY

Fosaprepitant, a prodrug of aprepitant, when administered intravenously is rapidly converted to aprepitant, a substance P/neurokinin 1 (NK₁) receptor antagonist. Plasma concentrations of fosaprepitant are below the limits of quantification (10 ng/mL) within 30 minutes of the completion of infusion.

Dexamethasone: Fosaprepitant 150 mg administered as a single intravenous dose on Day 1 increased the AUC_{0-24hr} of dexamethasone, a CYP3A4 substrate, by approximately 2.01, 1.86 and 1.18-fold on Days 1, 2 and 3 respectively when dexamethasone was coadministered as a single 8 mg oral dose on Days 1, 2, and 3. The oral dexamethasone dose on Days 1 and 2 should be reduced by approximately 50% when coadministered with fosaprepitant 150 mg IV on Day 1 to achieve exposures of dexamethasone similar to those obtained when given without fosaprepitant 150 mg.

Oral aprepitant, when given as a regimen of 125 mg with dexamethasone coadministered orally as 20 mg on Day 1, and oral aprepitant when given as 80 mg/day with dexamethasone coadministered orally as 8 mg on Days 2 through 5, increased the AUC of dexamethasone,

a CYP3A4 substrate by 2.2-fold, on Days 1 and 5. The daily dose of dexamethasone administered in clinical CINV studies with oral aprepitant reflects an approximate 50% reduction of the dose of dexamethasone (see DOSAGE AND ADMINISTRATION).

Methylprednisolone: Oral aprepitant, when given as a regimen of 125 mg on Day 1 and 80 mg/day on Days 2 and 3, increased the AUC of methylprednisolone, a CYP3A4 substrate, by 1.3-fold on Day 1 and by 2.5-fold on Day 3, when methylprednisolone was coadministered intravenously as 125 mg on Day 1 and orally as 40 mg on Days 2 and 3.

Warfarin: A single 125-mg dose of aprepitant was administered on Day 1 and 80 mg/day on Days 2 and 3 to healthy subjects who were stabilized on chronic warfarin therapy. Although there was no effect of oral aprepitant on the plasma AUC of R(+) or S(-) warfarin determined on Day 3, there was a 34% decrease in S(-) warfarin (a CYP2C9 substrate) trough concentration accompanied by a 14% decrease in the prothrombin time (reported as International Normalized Ratio or INR) 5 days after completion of dosing with oral aprepitant. In patients on chronic warfarin therapy, the prothrombin time (INR) should be closely monitored in the 2-week period, particularly at 7 to 10 days, following initiation of fosaprepitant with each chemotherapy cycle (see WARNINGS AND PRECAUTIONS).

Tolbutamide: Oral aprepitant, when given as 125 mg on Day 1 and 80 mg/day on Days 2 and 3, decreased the AUC of tolbutamide (a CYP2C9 substrate) by 23% on Day 4, 28% on Day 8, and 15% on Day 15, when a single dose of tolbutamide 500 mg was administered orally prior to the administration of the 3-day regimen of oral aprepitant and on Days 4, 8, and 15.

Oral contraceptives: Aprepitant, when given once daily for 14 days as a 100-mg capsule with an oral contraceptive containing 35 mcg of ethinyl estradiol and 1 mg of norethindrone, decreased the AUC of ethinyl estradiol by 43%, and decreased the AUC of norethindrone by 8%; therefore the efficacy of hormonal contraceptives during and for 28 days after administration of fosaprepitant may be reduced. Alternative or back-up methods of contraception should be used during treatment with fosaprepitant and for 1 month following administration (see WARNINGS AND PRECAUTIONS).

Midazolam: Fosaprepitant 150 mg administered as a single intravenous dose on Day 1 increased the AUC_{0-∞} of midazolam by approximately 1.8-fold on Day 1 and had no effect (1.0-fold) on Day 4 when midazolam was coadministered as a single oral dose of 2 mg on Days 1 and 4. Fosaprepitant 150 mg IV is a weak CYP3A4 inhibitor as a single dose on Day 1 with no evidence of inhibition or induction of CYP3A4 observed on Day 4.

In addition, when fosaprepitant was administered as a dose of 100 mg over 15 minutes along with a single dose of midazolam 2 mg, the plasma AUC of midazolam was increased by 1.6-fold.

Oral aprepitant increased the AUC of midazolam, a sensitive CYP3A4 substrate, by 2.3-fold on Day 1 and 3.3-fold on Day 5, when a single oral dose of midazolam 2 mg was coadministered on Day 1 and Day 5 of a regimen of oral aprepitant 125 mg on Day 1 and 80 mg/day on Days 2 through 5. The potential effects of increased plasma concentrations of midazolam or other

benzodiazepines metabolized via CYP3A4 (alprazolam, triazolam) should be considered when coadministering these agents with fosaprepitant or aprepitant.

In another study with intravenous administration of midazolam, oral aprepitant was given as 125 mg on Day 1 and 80 mg/day on Days 2 and 3, and midazolam 2 mg IV was given prior to the administration of the 3-day regimen of oral aprepitant and on Days 4, 8, and 15. Oral aprepitant increased the AUC of midazolam by 25% on Day 4 and decreased the AUC of midazolam by 19% on Day 8 relative to the dosing of oral aprepitant on Days 1 through 3. These effects were not considered clinically important. The AUC of midazolam on Day 15 was similar to that observed at baseline.

An additional study was completed with intravenous administration of midazolam and oral aprepitant. Intravenous midazolam 2 mg was given 1 hour after oral administration of a single dose of oral aprepitant 125 mg. The plasma AUC of midazolam was increased by 1.5-fold. Depending on clinical situations (e.g., elderly patients) and degree of monitoring available, dosage adjustment for intravenous midazolam may be necessary when it is coadministered with aprepitant for the chemotherapy induced nausea and vomiting indication (125 mg on Day 1 followed by 80 mg on Days 2 and 3).

Ketoconazole: When a single 125-mg dose of oral aprepitant was administered on Day 5 of a 10-day regimen of 400 mg/day of ketoconazole, a strong CYP3A4 inhibitor, the AUC of aprepitant increased approximately 5-fold and the mean terminal half-life of aprepitant increased approximately 3-fold. Concomitant administration of fosaprepitant with strong CYP3A4 inhibitors should be approached cautiously.

Rifampin: When a single 375-mg dose of oral aprepitant was administered on Day 9 of a 14-day regimen of 600 mg/day of rifampin, a strong CYP3A4 inducer, the AUC of aprepitant decreased approximately 11-fold and the mean terminal half-life decreased approximately 3-fold. Coadministration of fosaprepitant with drugs that induce CYP3A4 activity may result in reduced plasma concentrations and decreased efficacy.

Diltiazem: In a study in 10 patients with mild to moderate hypertension, intravenous infusion of 100 mg of fosaprepitant over 15 minutes with diltiazem 120 mg 3 times daily, resulted in a 1.5-fold increase of aprepitant AUC and a 1.4-fold increase in diltiazem AUC. It also resulted in a small but clinically meaningful further maximum decrease in diastolic blood pressure [mean (SD) of 24.3 (\pm 10.2) mm Hg with fosaprepitant versus 15.6 (\pm 4.1) mm Hg without fosaprepitant] and resulted in a small further maximum decrease in systolic blood pressure [mean (SD) of 29.5 (\pm 7.9) mm Hg with fosaprepitant versus 23.8 (\pm 4.8) mm Hg without fosaprepitant], which may be clinically meaningful, but did not result in a clinically meaningful further change in heart rate or PR interval, beyond those changes induced by diltiazem alone.

In patients with mild to moderate hypertension, administration of aprepitant once daily, as a tablet formulation comparable to 230 mg of the capsule formulation, with diltiazem 120 mg 3 times daily for 5 days, resulted in a 2-fold increase of aprepitant AUC and a simultaneous 1.7-fold

increase of diltiazem AUC. These pharmacokinetic effects did not result in clinically meaningful changes in ECG, heart rate, or blood pressure beyond those changes induced by diltiazem alone.

Paroxetine: Coadministration of once daily doses of aprepitant, as a tablet formulation comparable to 85 mg or 170 mg of the capsule formulation, with paroxetine 20 mg once daily, resulted in a decrease in AUC by approximately 25% and C_{max} by approximately 20% of both aprepitant and paroxetine.

TOXICOLOGY

Animal Toxicology

Acute Toxicity

The approximate oral LD_{50} of aprepitant was >2000 mg/kg in female mice and rats. The approximate intraperitoneal LD_{50} of aprepitant was >800 mg/kg, but <2000 mg/kg in female rats and >2000 mg/kg in female mice.

The approximate LD_{50} of fosaprepitant following intravenous administration was >500 mg/kg in female mice and >200 mg/kg in female rats

Chronic Toxicity

Fosaprepitant, when administered intravenously, is rapidly converted to aprepitant.

The toxicity potential of aprepitant was evaluated in a series of repeated-dose oral toxicity studies in rats and in dogs for up to 1 year.

In rats, oral administration of aprepitant for 6 months at doses up to the maximum feasible dose of 1000 mg/kg twice daily (approximately equivalent to [females] or lower than [males] the adult human dose based on systemic exposure following oral aprepitant 125 mg) produced increased hepatic weights that correlated with hepatocellular hypertrophy, increased thyroidal weights that correlated with thyroid follicular cell hypertrophy and/or hyperplasia, and pituitary cell vacuolation. These findings are a species-specific consequence of hepatic CYP enzyme induction in the rat, and are consistent with changes observed in rats with other structurally and pharmacologically dissimilar compounds that have been shown to induce hepatic CYP enzymes.

In dogs administered aprepitant orally for 9 months at doses ≥ 5 mg/kg twice daily (greater than or equal to 13 times the adult human dose based on systemic exposure following oral aprepitant 125 mg), toxicity was characterized by slight increases in serum alkaline phosphatase activity and decreases in the albumin/globulin ratio. Significantly decreased body weight gain, testicular degeneration, and prostatic atrophy were observed at doses ≥ 25 mg/kg twice daily (greater than or equal to 31 times the adult human dose based on systemic exposure following oral aprepitant 125 mg). A slight increase in hepatic weights with no histologic correlate was seen at 500 mg/kg twice daily (70 times the adult human dose based on systemic exposure following oral aprepitant

125 mg). No toxicity was observed in dogs administered 32 mg/kg/day (6 times the adult human dose based on systemic exposure following oral aprepitant 125 mg) for 1 year.

Local Tolerance

In rabbits, EMEND® IV caused initial focal acute inflammation when administered intravenously, paravenously, and subcutaneously. Focal skeletal muscle degeneration and necrosis with associated neutrophilic inflammation were noted with intramuscular injection. At the end of the follow-up period (post-dose day 8), paravenous injection sites showed focal subacute inflammation. Intramuscular injection site changes consisted of focal skeletal muscle necrosis and mineralization bordered by subacute inflammation and focal skeletal muscle regeneration.

Carcinogenesis

Carcinogenicity studies were conducted in mice and rats for approximately 2 years with oral aprepitant. In mice, aprepitant was not carcinogenic at doses up to 500 mg/kg/day (approximately 2 times the adult human dose based on systemic exposure). Rats developed hepatocellular adenomas at a dose of 25 mg/kg twice daily (females) and 125 mg/kg twice daily (females and males), thyroid follicular cell adenomas at a dose of 125 mg/kg twice daily (females and males), and thyroid follicular cell carcinomas at a dose of 125 mg/kg twice daily (males). Systemic exposures at these doses in rats were approximately equivalent to or lower than exposures in humans at the recommended dose. Tumors of these types are a species-specific consequence of hepatic CYP enzyme induction in the rat, and are consistent with changes observed in rats with other structurally and pharmacologically dissimilar compounds that have been shown to induce hepatic CYP enzymes. Carcinogenicity studies were not conducted with fosaprepitant.

Mutagenesis

Fosaprepitant and aprepitant were neither mutagenic nor genotoxic in assays conducted to detect mutagenicity, DNA strand breaks, and chromosomal aberrations. Aprepitant was negative in the *in vitro* microbial and TK6 human lymphoblastoid cell mutagenesis assays, the *in vitro* alkaline elution/rat hepatocyte DNA strand break test, the *in vitro* chromosomal aberration assay in Chinese hamster ovary cells, and the *in vivo* mouse micronucleus assay in bone marrow.

Reproduction

Fosaprepitant, when administered intravenously, is rapidly converted to aprepitant. In the fertility studies conducted with fosaprepitant and aprepitant, the highest systemic exposures to aprepitant were obtained following oral administration of aprepitant.

Aprepitant administered to female rats at doses up to the maximum feasible dose of 1000 mg/kg twice daily (approximately equivalent to the adult human dose based on systemic exposure following oral aprepitant 125 mg) had no effects on mating performance, fertility, or embryonic/fetal survival.

Administration of aprepitant to male rats at doses up to the maximum feasible dose of 1000 mg/kg twice daily (lower than the adult human dose based on systemic exposure following oral aprepitant 125 mg) produced no effects on mating performance, fertility, embryonic/fetal survival, sperm count and motility, testicular weights, or the microscopic appearance of the testes and epididymides.

Development

Fosaprepitant, when administered intravenously, is rapidly converted to aprepitant. In the teratology studies conducted with fosaprepitant and aprepitant, the highest systemic exposures to aprepitant were obtained following oral administration of aprepitant.

In rats and rabbits administered oral doses of aprepitant up to 1000 mg/kg twice daily and 25 mg/kg/day, respectively (up to 1.5 times the systemic exposure at the adult human dose following oral aprepitant 125 mg), there was no evidence of developmental toxicity as assessed by embryonic/fetal survival, fetal body weight, and fetal external, visceral, and skeletal morphology. Placental transfer of aprepitant occurred in rats and rabbits at these doses. Concentrations of aprepitant in fetal plasma were approximately 27% and 56% of maternal plasma concentrations in rats and rabbits, respectively.

Significant concentrations of aprepitant were observed in the milk of lactating rats administered 1000 mg/kg twice daily. At this dose, the mean milk drug concentration was 90% of the mean maternal plasma concentration.

REFERENCES

1. MRL Clinical Study Report: A Randomized, Double-Blind, Placebo-Controlled, Parallel-Group Study, Conducted Under In-House Blinding Conditions, to Examine the Safety, Tolerability, and Efficacy of MK-0869 for the Prevention of Chemotherapy-Induced Nausea and Vomiting Associated With High-Dose Cisplatin (Protocol 052).
2. MRL Clinical Study Report: A Randomized, Double-Blind, Placebo-Controlled, Parallel-Group Study, Conducted Under In-House Blinding Conditions, to Examine the Safety, Tolerability, and Efficacy of MK-0869 for the Prevention of Chemotherapy-Induced Nausea and Vomiting Associated With High-Dose Cisplatin (Protocol 054).
3. MRL Clinical Study Report, Multicenter Study: A Randomized, Double-Blind, Parallel-Group Study Conducted Under In-House Blinding Conditions to Determine the Efficacy and Tolerability of Aprepitant for the Prevention of Chemotherapy-Induced Nausea and Vomiting Associated With Moderately Emetogenic Chemotherapy (Protocol 071).
4. MRL Clinical Study Report, A Randomized, Double-Blind, Parallel-Group Study Conducted Under In-House Blinding Conditions to Determine the Efficacy and Tolerability of Aprepitant for the Prevention of Chemotherapy-Induced Nausea and Vomiting Associated With Moderately Emetogenic Chemotherapy (Protocol 130).
5. MRL Clinical Study Report, Multicenter Study: A Randomized, Double-Blind, Parallel-Group Study Conducted Under In-House Blinding Conditions to Examine the Safety, Tolerability, and Efficacy of a Single Dose of MK-0517 for the Prevention of Chemotherapy-Induced Nausea and Vomiting (CINV) Associated With Cisplatin Chemotherapy (Protocol 017L).
6. de Wit R, Herrstedt J, Rapoport B, Carides AD, Guoguang-Ma J, Elmer M et al. The oral NK(1) antagonist, aprepitant, given with standard antiemetics provides protection against nausea and vomiting over multiple cycles of cisplatin-based chemotherapy: a combined analysis of two randomised, placebo-controlled phase III clinical trials. *Eur J Cancer* 2004;40:403–10.
7. Gralla RJ, de Wit R, Herrstedt J, Carides AD, Ianus J, Guoguang-Ma J et al. Antiemetic Efficacy of the Neurokinin-1 Antagonist, Aprepitant, Plus a 5HT3 Antagonist and a Corticosteroid in Patients Receiving Anthracyclines or Cyclophosphamide in Addition to High-Dose Cisplatin. *Cancer* 2005 Aug 15;104(4):864–8.
8. Hesketh PJ, Grunberg SM, Gralla RJ, Warr DG, Roila F, de Wit R et al. The oral neurokinin-1 antagonist aprepitant for the prevention of chemotherapy-induced nausea and vomiting: a multinational, randomized, double-blind, placebo-controlled trial in patients receiving high-dose cisplatin—the Aprepitant Protocol 052 Study Group. *J Clin Oncol* 2003;21:4112–9.

9. Hesketh PJ, Grunberg, SM, Herrstedt J, de Wit R, Gralla RJ, Carides AD. Combined data from two phase III trials of the NK1 antagonist aprepitant plus a 5HT3 antagonist and a corticosteroid for prevention of chemotherapy-induced nausea and vomiting: effect of gender on treatment response. *Support Care Cancer* 2006;14:354–60.
10. Poli-Bigelli S, Rodrigues-Pereira J, Carides AD, Guoguang-Ma J, Eldridge K, Hipple A et al. Addition of the neurokinin 1 receptor antagonist aprepitant to standard antiemetic therapy improves control of chemotherapy-induced nausea and vomiting. Results from a randomized, double-blind, placebo-controlled trial in Latin America. *Cancer* 2003;97:3090–8.
11. Rapoport BL, Jordan K, Boice JA, Taylor A, Brown C, Hardwick JS et al. Aprepitant for the prevention of chemotherapy-induced nausea and vomiting associated with a broad range of moderately emetogenic chemotherapies and tumor types: a randomized, double blind study. *Support Care Cancer* 2009 Jul 01; published on line.
12. Warr DG, Hesketh PJ, Gralla RJ, Muss HB, Herrstedt J, Eisenbert PD et al. Efficacy and tolerability of aprepitant for the prevention of chemotherapy-induced nausea and vomiting in patients with breast cancer after one cycle of moderately emetogenic chemotherapy. *J Clin Oncol* 2005;23:2822–30.

PART III: CONSUMER INFORMATION

EMEND® IV
fosaprepitant for injection
(as fosaprepitant dimeglumine)

This leaflet is part III of a three-part "Product Monograph" published when EMEND® IV were approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about EMEND® IV. Contact your physician or pharmacist if you have any questions about the drug.

Please read this leaflet carefully before you start to take your medicine, even if you have just refilled your prescription. Some of the information in the previous leaflet may have changed.

Remember that your physician has prescribed this medicine only for you. Never give it to anyone else.

ABOUT THIS MEDICATION**What the medication is used for:**

EMEND® IV, in combination with 5-HT₃ antagonists and dexamethasone, is indicated for the prevention of nausea and vomiting associated with your cancer chemotherapy treatment.

What it does:

EMEND® IV is a member of a class of medicines called neurokinin 1 (NK₁) receptor antagonists. Fosaprepitant when administered intravenously is rapidly converted to aprepitant. Aprepitant works by blocking neurokinin, a substance in the brain that causes nausea and vomiting.

When it should not be used:

Do not take EMEND® IV if you are allergic to fosaprepitant dimeglumine, aprepitant or any of the other ingredients of EMEND® IV.

Do not take EMEND® IV with pimozide, terfenadine, astemizole, or cisapride. Taking EMEND® IV with these medications could result in **serious or life-threatening problems**.

What the medicinal ingredient is:

Fosaprepitant dimeglumine

What the important non-medicinal ingredients are:

edetate disodium, polysorbate 80, lactose anhydrous, sodium hydroxide and/or hydrochloric acid (for pH adjustment).

What dosage forms it comes in:

Powder for injection. Each vial contains 150 mg of fosaprepitant.

WARNINGS AND PRECAUTIONS**Serious Warnings and Precautions**

Drug interactions with:

- Medicines that are likely to be broken down mainly by the liver
- Warfarin
- Hormonal contraception (birth control medicines)

Possible serious side effects:

- Severe allergic (hypersensitivity) reactions during infusion

BEFORE you use EMEND® IV talk to your physician or pharmacist if:

- you have any past or present medical problems
- you have liver problems
- you have any allergies
- you drive a car or operate machinery
- you are pregnant or plan to become pregnant
- you are breast-feeding or plan to breast-feed

Use in children

EMEND® IV should not be given to children under 18 years of age.

Use in the elderly

No dosage adjustment is necessary.

INTERACTIONS WITH THIS MEDICATION

Tell your physician about all medicines that you are taking or plan to take, even those you can get without a prescription or herbal products.

Your physician may check that your medicines are working properly together if you are taking other medicines such as:

- anti-anxiety drugs (such as alprazolam, midazolam)
- birth control medicines (which may not work as well)
- ketoconazole (an antifungal)
- rifampin (an antibiotic)
- paroxetine (a medicine used to treat a certain type of depression)
- diltiazem (a medicine used to treat high blood pressure)
- dexamethasone, methylprednisolone (steroid medicines used for a variety of conditions)
- warfarin (a blood thinner)
- tolbutamide (a medicine used to treat diabetes)
- phenytoin (a medicine used to treat seizures)

Following the infusion of EMEND® IV, the early concentration of aprepitant in the blood is double that of an EMEND® capsule (125 mg); therefore, a possible risk for increased side effects cannot be ruled out.

PROPER USE OF THIS MEDICATION

EMEND[®] IV may be taken with or without food.

Usual dose:

EMEND[®] IV 150 mg given on Day 1 only.

- Day of chemotherapy: EMEND[®] IV 150 mg will be given to you intravenously approximately 30 minutes before you start your chemotherapy treatment.

Overdose:

In case of a drug overdose, contact a health care practitioner, hospital emergency department or Regional Poison Control Centre, immediately.

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

Any medicine may have unintended or undesirable effects, so-called side effects.

Like all prescription drugs, EMEND[®] IV may cause side effects. The most common side effects included diarrhea, stomach pain, upset stomach, vomiting, dizziness, hiccups, fatigue, weakness, constipation, headache, and loss of appetite.

Infusion-site reactions included pain, hardening, swelling of a vein caused by a blood clot, redness, and/or itching at the infusion site.

Other side effects may also occur rarely, which include: anxiousness, fever with increased risk of infection, dry mouth, conjunctivitis (eye discharge and itching), excessive sweating, flushing, painful burning urination, muscle cramp or pain, taste disturbance, high blood pressure, ringing in the ear (tinnitus) and low blood pressure.

Allergic reactions, which may be sudden and/or serious, and may include hives, rash, itching, redness of the face/skin, and cause difficulty in breathing or swallowing.

Ask your physician or pharmacist for more information. Both have a more complete list of side effects. Tell your physician or pharmacist promptly about these or any other unusual symptoms.

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

Symptoms / Effects		Talk with your physician or pharmacist		Stop taking drug and call your physician or pharmacist
		Only if severe	In all cases	
Uncommon	Allergic reactions/ Angioedema (swelling of the face, eyes, lips, tongue, throat, difficulty in breathing or swallowing)		√	
Uncommon	Stevens-Johnson syndrome/toxic epidermal necrolysis (severe skin reactions, blistering)		√	
Uncommon	Urticaria (severe rash, itching, swelling of the hands and feet)		√	

This is not a complete list of side effects. For any unexpected effects while taking EMEND[®] IV, contact your physician or pharmacist.

HOW TO STORE IT

Vials: Sterile powder for intravenous use. Store at 2–8°C.

Keep EMEND[®] IV and all medicines safely away from children.

REPORTING SUSPECTED SIDE EFFECTS

You can report any suspected adverse reactions associated with the use of health products to the Canada Vigilance Program by one of the following 3 ways:

- Report online at www.healthcanada.gc.ca/medeffect
- Call toll-free at 1-866-234-2345
- Complete a Canada Vigilance Reporting Form and:
 - Fax toll-free to 1-866-678-6789, or
 - Mail to: Canada Vigilance Program
Health Canada
Postal Locator 0701E
Ottawa, ON K1A 0K9

Postage paid labels, Canada Vigilance Reporting Form and the adverse reaction reporting guidelines are available on the MedEffect[™] Canada Web site at www.healthcanada.gc.ca/medeffect

NOTE: Should you require information related to the management of side effects, contact your health professional. The Canada Vigilance Program does not provide medical advice.

MORE INFORMATION

Also, you can report any suspected adverse reactions associated with the use of health products to Merck Canada Inc. by one of the following 2 ways:

- Call toll-free at 1-800-567-2594
- Complete a Canada Vigilance Reporting Form and:
 - Fax toll-free to 1-800-369-3090, or
 - Mail to: Merck Canada Inc.
Pharmacovigilance
P.O. Box 1005
Pointe-Claire-Dorval, QC H9R 4P8

This document plus the full product monograph, prepared for health professionals can be found at:

<http://www.merck.ca>

or by contacting the sponsor, Merck Canada Inc.

at: 1-800-567-2594

This leaflet was prepared by Merck Canada Inc.

Last revised: January 30, 2012

® Registered trademark of Merck Sharp & Dohme Corp., a subsidiary of **Merck & Co., Inc.** Used under license.

© 2011, Merck Canada Inc., a subsidiary of **Merck & Co., Inc.**
All rights reserved.

