

## PRODUCT MONOGRAPH

Pr **ZENHALE™**

mometasone furoate / formoterol fumarate dihydrate

Inhalation aerosol

Pr ZENHALE™ 50/5

50 mcg mometasone furoate / 5 mcg formoterol fumarate dihydrate

Pr ZENHALE™ 100/5

100 mcg mometasone furoate / 5 mcg formoterol fumarate dihydrate

Pr ZENHALE™ 200/5

200 mcg mometasone furoate / 5 mcg formoterol fumarate dihydrate

Corticosteroid and long-acting beta<sub>2</sub>-agonist combination for oral inhalation

**Merck Canada Inc.**  
Kirkland, Québec  
H9H 4M7

Date of Preparation:  
January 7, 2011  
Date of Revision:  
June 30, 2011

Submission Control No: 147075

## Table of Contents

<b>PART I: HEALTH PROFESSIONAL INFORMATION.....</b>	<b>3</b>
SUMMARY PRODUCT INFORMATION .....	3
INDICATIONS AND CLINICAL USE.....	3
CONTRAINDICATIONS .....	4
WARNINGS AND PRECAUTIONS.....	4
ADVERSE REACTIONS.....	9
DRUG INTERACTIONS .....	11
DOSAGE AND ADMINISTRATION .....	12
OVERDOSAGE .....	15
ACTION AND CLINICAL PHARMACOLOGY .....	15
STORAGE AND STABILITY.....	19
SPECIAL HANDLING INSTRUCTIONS .....	19
DOSAGE FORMS, COMPOSITION AND PACKAGING .....	19
<b>PART II: SCIENTIFIC INFORMATION.....</b>	<b>20</b>
PHARMACEUTICAL INFORMATION.....	20
CLINICAL TRIALS.....	21
DETAILED PHARMACOLOGY .....	26
MICROBIOLOGY .....	28
TOXICOLOGY .....	28
REFERENCES .....	31
<b>PART III: CONSUMER INFORMATION.....</b>	<b>32</b>

Pr **ZENHALE™**

mometasone furoate / formoterol fumarate dihydrate inhalation aerosol

**PART I: HEALTH PROFESSIONAL INFORMATION**

**SUMMARY PRODUCT INFORMATION**

<b>Route of Administration</b>	<b>Dosage Form / Strength</b>	<b>Nonmedicinal Ingredients</b>
oral	Inhalation aerosol / 50, 100 or 200 mcg mometasone furoate & 5 mcg formoterol fumarate dihydrate per actuation	Ethanol, HFA (propellant), oleic acid

**INDICATIONS AND CLINICAL USE**

ZENHALE™ (mometasone furoate / formoterol fumarate dihydrate inhalation aerosol), is indicated for the treatment of asthma, in patients 12 years of age and older with reversible obstructive airway disease.

ZENHALE™ is **not** indicated for patients whose asthma can be managed by occasional use of a rapid onset, short duration, inhaled beta<sub>2</sub>-agonist, or for patients whose asthma can be successfully managed by inhaled corticosteroids along with occasional use of a rapid onset, short duration, inhaled beta<sub>2</sub>-agonist.

Long-acting beta<sub>2</sub>-adrenergic agonists (LABA), such as formoterol, one of the active ingredients in ZENHALE™, may increase the risk of asthma-related death. Available data from controlled clinical trials suggest that LABA increase the risk of asthma-related hospitalization in pediatric and adolescent patients (see WARNINGS AND PRECAUTIONS). Therefore, when treating patients with asthma, physicians should only prescribe ZENHALE™ for patients not adequately controlled on a long-term asthma control medication, such as an inhaled corticosteroid or whose disease severity clearly warrants initiation of treatment with both an inhaled corticosteroid and LABA.

Once asthma control has been achieved and maintained, assess the patient at regular intervals and do not use ZENHALE™ for patients whose asthma can be adequately controlled on low or medium dose inhaled corticosteroids.

**Geriatrics (≥65 years of age):**

Based on available data for ZENHALE™ or its active components, no adjustment of ZENHALE™ in geriatric patients is warranted.

**Pediatrics (<12 years of age):**

The safety and efficacy of ZENHALE™ have not been established in children less than 12 years of age.

**CONTRAINDICATIONS**

- Patients who are hypersensitive to this drug or to any ingredient in the formulation or component of the container. For a complete listing, see the DOSAGE FORMS, COMPOSITION AND PACKAGING section of the product monograph.

**WARNINGS AND PRECAUTIONS****WARNING**

**Asthma-Related Death:** Long-acting  $\beta_2$ -adrenergic agonists (LABA), such as formoterol, one of the active ingredients in ZENHALE™, may increase the risk of asthma-related death. Data from a large placebo-controlled US study, which compared the safety of salmeterol, another LABA, or placebo added to patients' usual asthma therapy showed an increase in asthma-related deaths in patients receiving salmeterol. This finding with salmeterol is considered a class effect of the LABA, including formoterol. Available data from controlled clinical trials suggest that LABA increase the risk of asthma-related hospitalization in pediatric and adolescent patients. Currently available data are inadequate to determine whether concurrent use of inhaled corticosteroids or other long-term asthma control drugs mitigates the increased risk of asthma-related death from LABA. Therefore, when treating patients with asthma, ZENHALE™ should only be used for patients not adequately controlled on a long-term asthma control medication, such as an inhaled corticosteroid or whose disease severity clearly warrants initiation of treatment with both an inhaled corticosteroid and LABA. Once asthma control is achieved and maintained, assess the patient at regular intervals and do not use ZENHALE™ for patients whose asthma can be adequately controlled on low or medium dose inhaled corticosteroids (see DOSAGE AND ADMINISTRATION).

**General**

ZENHALE™ should not be used to treat acute symptoms of asthma. Adequate education should be provided to the patient regarding the use of long-acting beta<sub>2</sub>-agonists and the acute treatment of asthma, with close-follow up to ensure compliance (see **Respiratory**).

**Discontinuance:** Treatment with inhaled corticosteroids should not be stopped abruptly in patients with asthma due to risk of exacerbation. In this case, therapy should be titrated down gradually, under physician supervision.

### ***Transferring from Systemic Corticosteroid Therapy***

Particular care is needed for patients who are transferred from systemically active corticosteroids to ZENHALE™, because deaths due to adrenal insufficiency have occurred in asthmatic patients during and after transfer from systemic corticosteroids to less systemically available inhaled corticosteroids. After withdrawal from systemic corticosteroids, a number of months are required for recovery of hypothalamic-pituitary-adrenal (HPA) axis function.

During periods of stress, including trauma, surgery, or infection, or a severe asthma attack, patients transferred from systemic corticosteroids will require supplementary treatment with a short course of systemic corticosteroids, which is gradually tapered as symptoms subside. It is recommended that such patients carry a supply of oral corticosteroids and a warning card indicating their need and recommended dosage of systemic corticosteroids during stressful periods. Periodic testing of adrenocortical function, particularly measurement of early morning plasma cortisol levels, is recommended.

Transfer of patients from systemic corticosteroid therapy to ZENHALE™ may unmask pre-existing allergic conditions previously suppressed by systemic corticosteroid therapy. If this occurs, symptomatic treatment is recommended.

### ***Systemic Effects of Corticosteroids***

Systemic effects of inhaled corticosteroids may occur, particularly at high doses prescribed for prolonged periods. These effects are much less likely to occur than with oral corticosteroids. Possible systemic effects include Cushing's syndrome, Cushingoid features, adrenal suppression (see **Endocrine and Metabolism**), growth retardation in children and adolescents, decrease in bone mineral density, cataracts and glaucoma (see **Ophthalmologic**). Therefore, it is important that the dose of ZENHALE™ is titrated to the lowest dose at which effective control of asthma is maintained.

### ***Other Long-Acting Beta<sub>2</sub>-agonist***

ZENHALE™ should not be used in conjunction with another long-acting beta<sub>2</sub>-agonist.

### ***Exceeding Recommended Dose***

**Do not exceed recommended dosage with ZENHALE™.**

The dose of ZENHALE™ should be individualized to the patient's needs and should be at the lowest possible dose to fulfill the therapeutic objective. It should not be increased beyond the maximum recommended dose (see DOSAGE AND ADMINISTRATION). No evidence supports that the administration of ZENHALE™ in amounts greater than recommended doses increases efficacy.

## **Cardiovascular**

Although clinically not significant, a small increase in QTc interval has been reported with therapeutic doses of formoterol. It is not known if this becomes clinically significant when concomitant medications causing similar effects are prescribed and/or in the presence of heart diseases, hypokalemia, or hypoxia.

ZENHALE™, like other beta<sub>2</sub>-agonist containing products, should be used with caution in patients with ischemic heart disease, cardiac arrhythmias (especially third-degree atrioventricular block), severe cardiac decompensation, idiopathic subvalvular aortic stenosis, hypertrophic obstructive cardiomyopathy, thyrotoxicosis, known or suspected prolongation of the QT interval (QTc >0.44 sec).

## **Dependence/Tolerance**

There is no information to indicate that abuse or dependency occur with mometasone furoate or formoterol fumarate.

## **Ear/Nose/Throat**

### ***Oropharyngeal candidiasis***

During clinical trials with ZENHALE™, oral candidiasis, which is associated with the use of inhaled glucocorticosteroids, occurred in some patients. This infection may require treatment with appropriate antifungal therapy and in some patients discontinuance of ZENHALE™ may be necessary. Patients should be advised to rinse their mouth after inhalation of ZENHALE™.

## **Endocrine and Metabolism**

### ***Adrenal Suppression***

ZENHALE™ will usually permit control of asthma symptoms with less suppression of HPA axis function than therapeutically equivalent oral doses of prednisone.

When using inhaled corticosteroids, the possibility for clinically significant adrenal suppression may occur, especially after treatment with higher than recommended doses. This must be considered during periods of stress or elective surgery, when additional systemic corticosteroids may be needed. During clinical trials the effects of ZENHALE™ (at doses of mometasone furoate 800 mcg/day) on plasma cortisol were not clinically significant.

### ***Hypokalemia and Hyperglycemia***

Potentially serious hypokalemia may occur as a result of beta<sub>2</sub>-agonist therapy. Hypokalemia may increase susceptibility to cardiac arrhythmias. Particular caution is advised in patients with severe asthma as hypokalemia may be potentiated by hypoxia and concomitant treatment (see DRUG INTERACTIONS / Drug-Drug Interactions). It is recommended that serum potassium levels be monitored in such situations.

Due to the hyperglycemic effect of beta<sub>2</sub>-stimulants, including formoterol, additional blood glucose monitoring is recommended in diabetic patients.

There is an enhanced effect of corticosteroids on patients with hypothyroidism.

### **Hepatic/Biliary/Pancreatic**

There is an enhanced effect of corticosteroids on patients with cirrhosis.

### **Immune**

Use ZENHALE™ with caution, if at all, in patients with active or quiescent tuberculosis infections of the respiratory tract, or in untreated systemic fungal, bacterial, systemic viral infections or ocular herpes simplex.

Advise patients who are receiving corticosteroids or other immunosuppressant medicines of the risk of exposure to certain infections (e.g. chickenpox, measles) and of the importance of obtaining medical advice if such exposure occurs. This is of particular importance in children. If exposed to chickenpox, prophylaxis with varicella zoster immune globulin (VZIG) may be indicated. If exposed to measles, prophylaxis with pooled intravenous immunoglobulin (IG) may be indicated. If chickenpox develops, treatment with antiviral agents may be considered.

Immediate hypersensitivity, reactions including angioedema, urticaria, flushing and bronchospasm may occur after administration of ZENHALE™.

### **Monitoring and Laboratory Tests**

#### ***Monitoring Control of Asthma***

ZENHALE™ should not be introduced in acutely deteriorating asthma, which is a potentially life threatening condition. Increasing use of rapid onset, short duration inhaled bronchodilators to control symptoms indicates deterioration of asthma control. Sudden and progressive deterioration in asthma control is potentially life-threatening and the treatment plan should be re-evaluated. Also, where current dosage of ZENHALE™ has failed to give adequate control of reversible obstructive airways disease the patient should be reviewed by a physician (see DOSAGE AND ADMINISTRATION).

Before introducing ZENHALE™, adequate education should be provided to the patient on how to use the drug and what to do if asthma flares up.

During long-term therapy, HPA axis function and haematological status should be assessed periodically. For patients at risk, monitoring of bone and ocular effects (cataract and glaucoma) should also be considered in patients receiving maintenance therapy with ZENHALE™. It is recommended that the height of children and adolescents receiving prolonged treatment with inhaled corticosteroids is regularly monitored.

### **Ophthalmologic**

Rare cases of cataracts and glaucoma have been reported with use of mometasone furoate. For patients at risk, monitoring of ocular effects (cataract and glaucoma) should be considered in patients receiving maintenance therapy with ZENHALE™.

## **Respiratory**

### ***Acute Asthma Episodes***

ZENHALE™ is not indicated for rapid relief of bronchospasm or other acute episodes of asthma. In the event of an acute attack, a short-acting  $\beta_2$ -agonist should be used. A short-acting  $\beta_2$ -agonist should be available to the patient at all times. Patients must be informed of the need to seek medical treatment immediately if their asthma deteriorates suddenly.

### ***Asthma exacerbations***

Serious asthma-related adverse events and exacerbations may occur during treatment with ZENHALE™, like other beta<sub>2</sub>-agonist containing products. Patients should be asked to continue treatment but to seek medical advice if asthma symptoms remain uncontrolled or worsen after initiation of ZENHALE™.

ZENHALE™ should not be initiated in patients during rapidly deteriorating or potentially-life threatening episodes of asthma. ZENHALE™ has not been studied in patients with acutely deteriorating asthma.

The prescriber should reassess asthma therapy if symptoms persist; dosing requirements are increased to maintain control, asthmatic episodes not responsive to bronchodilators, or decreased peak flow, because this usually indicates that the underlying condition has deteriorated. During such episodes, consideration should be given to additional corticosteroid therapies.

### ***Inhalation induced bronchospasm***

As with other inhalation therapy, the potential for inhalation induced bronchospasm should be kept in mind. If it occurs, the preparation should be discontinued immediately and alternative therapy substituted.

## **Special Populations**

**Pregnant Women:** There are no adequate and well-controlled studies of ZENHALE™ use in pregnant women. Studies in animals with mometasone furoate, like other glucocorticoids, have shown reproductive toxicity (see TOXICOLOGY); however the potential risk for humans is unknown. ZENHALE™ should not be used during pregnancy unless the potential benefit justifies the potential risk to the fetus.

Infants born of mothers who received corticosteroids during pregnancy are to be observed carefully for hypoadrenalism.

**Use in Labour and Delivery:** Like other beta<sub>2</sub>-adrenergic stimulants, formoterol may inhibit labour due to a relaxant effect on uterine smooth muscle.

**Nursing Women:** There are no data from well-controlled human studies on the use of ZENHALE™ in nursing mothers. Glucocorticoids are excreted in human milk. The excretion of mometasone furoate into human breast milk has not been investigated. Formoterol has been detected in the milk of rats. Based on data for the individual components, a decision on whether to continue/discontinue therapy with ZENHALE™ should be made taking into account the benefit of breast-feeding to the child and the benefit of ZENHALE™ therapy to the woman.

**Pediatrics:** The safety and efficacy of ZENHALE™ have not been established in children less than 12 years of age.

**Geriatrics (>65 years of age):** No overall differences in safety and effectiveness were observed between these patients and younger patients, in clinical studies. There is no adjustment of dosage of ZENHALE™ in geriatric patients.

## **ADVERSE REACTIONS**

### **Adverse Drug Reaction Overview**

ZENHALE™ contains both mometasone furoate and formoterol, therefore the type and severity of adverse reactions associated with each individual component of ZENHALE™ may be expected. There is no evidence of additional adverse events following concurrent administration of the two components.

Long-acting beta<sub>2</sub>-adrenergic agonists (LABA), such as formoterol, one of the active ingredients in ZENHALE™, may increase the risk of asthma-related death. Currently available data are inadequate to determine whether concurrent use of inhaled corticosteroids mitigates the increased risk of asthma-related death from LABA. Available data from controlled clinical trials suggest that LABA increase the risk of asthma-related hospitalization in pediatric and adolescent patients (see WARNINGS AND PRECAUTIONS).

Tremor, palpitations, electrocardiogram QT prolongation, tachycardia, hypertension and headache have been reported and are associated with pharmacological side effects of beta<sub>2</sub>-agonist treatment (including ZENHALE™). Cardiac arrhythmias (including atrial fibrillation, supraventricular tachycardia and ventricular extrasystoles) may occur in some patients.

Rarely, hypersensitivity reactions, including rash, urticaria, bronchospasm, arthralgia, angioedema, and anaphylactic reaction may occur in some patients.

Due to the mometasone furoate component for oral inhalation, oral candidiasis can occur in some patients. Incidence of oral candidiasis may be reduced by rinsing the mouth with water after using the product. Symptomatic candidiasis can be treated with topical antifungal therapy.

Systemic and local corticosteroid use may also result in the following:

- Immunosuppression
- Hypercorticism and adrenal suppression
- Growth retardation in children
- Glaucoma and cataracts
- Reduced bone density, osteoporosis and fracture

As with other inhalation therapy, inhalation induced bronchospasm may occur rarely.

See WARNINGS AND PRECAUTIONS for more detailed information.

## **Clinical Trial Adverse Drug Reactions**

*Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.*

Safety data is based on the 3 pivotal clinical trials (P04073, P04334, and P04431) and the long-term safety trial (P04139). The total number of patients (12 years of age and older with asthma) participating in these studies was 2659, of which 1131 were exposed to ZENHALE™. Eight hundred and sixty (860) patients were exposed to ZENHALE™ in the 12 to 26 week studies and 271 patients were exposed to ZENHALE™ in the 1 year study.

Table 1 demonstrates the incidence of treatment related adverse reactions associated with ZENHALE™ based upon the pooled data of the three pivotal clinical trials.

**Table 1: Treatment related adverse reactions in ZENHALE™ groups occurring at an incidence of ≥1% and more commonly than placebo**

Adverse Reactions	ZENHALE™*			Mometasone Furoate*			Formoterol*	Placebo*
	50/5 n=182 n (%)	100/5 n=424 n (%)	200/5 n=255 n (%)	50 mcg n=188 n (%)	100 mcg n=192 n (%)	200 mcg n=240 n (%)	5 mcg n=390 n (%)	n=384 n (%)
Oral Candidiasis	3 (1.6)	4 (0.9)	4 (1.6)	1 (0.5)	1 (0.5)	2 (0.8)	3 (0.8)	3 (0.8)
Nausea	2 (1.1)	0	0	0	0	0	0	0
Headache	2 (1.1)	1 (0.2)	1 (0.4)	1 (0.5)	1 (0.5)	0	2 (0.5)	2 (0.5)
Pharyngolaryngeal pain	2 (1.1)	0	0	1 (0.5)	0	0	3 (0.8)	0
Average Duration of Exposure (days)	162	116	81	159	165	79	139	137

These results are based on clinical trials P04073, P04334 & P04431.

\*All treatments were administered as two inhalations twice daily.

In a comparator safety study of one year treatment duration, patients 12 years of age and older were treated with medium dose ZENHALE™ 100/5 (n=141), high dose ZENHALE™ 200/5 (n=130) or an active comparator (n=133, 68 medium dose and 65 high dose inhaled corticosteroid/LABA combination).

Safety outcomes were similar to those observed in the 12 to 26 week trials and no treatment-related deaths or clinically judged asthma deteriorations or reduction in lung function were observed.

Dysphonia was observed at a higher frequency in the longer term treatment trial at a reported incidence of 7/141 (5%) patients receiving ZENHALE™ 100/5 and 4/130 (3.1%) patients receiving ZENHALE™ 200/5. Overall, through 52 weeks of observation, 15 patients demonstrated a ≥1.0 point change in LOCS III score (measured at the Week 26 and Week 52 timepoints using the Lens Opacities Classification System, Version III) from Baseline. At

Week 26, in the medium dose group, 2 (1.4%) patients receiving ZENHALE™ 100/5 and 4 (5.9%) patients receiving an active comparator demonstrated ocular changes. In the high dose group, 3 (2.3%) patients receiving ZENHALE™ 200/5 demonstrated ocular changes (no patients in the active comparator group). At Week 52, in the medium dose group, 4 (2.8%) patients receiving ZENHALE™ 100/5 and 1 (1.5%) patient receiving an active comparator demonstrated ocular changes. In the high dose group 3 (2.3%) patients receiving ZENHALE™ 200/5 and 1 (1.5%) patient receiving an active comparator demonstrated ocular changes. No incidences of appearance of posterior subcapsular cataracts typically associated with chronic use of high dose inhaled corticosteroid were reported in this clinical study. No clinically significant changes in blood chemistry, hematology, or ECG were observed.

### **Less Common Clinical Trial Adverse Drug Reactions (<1%)**

The following additional treatment related adverse reactions occurred in these clinical trials (P04073, P04334, P04431, P04139) in patients using ZENHALE™ with an incidence of <1% and occurred at a greater incidence than placebo:

*Cardiac Disorders:* tachycardia, palpitations

*Gastrointestinal Disorders:* dry mouth

*Immune System Disorders:* hypersensitivity reactions with the following manifestations – bronchospasm, dermatitis allergic, urticaria

*Infections and Infestations:* pharyngitis

*Musculoskeletal and Connective Tissue Disorders:* muscle spasms\*

*Nervous System Disorders:* tremor, dizziness\*

*Psychiatric Disorders:* insomnia, nervousness\*

*Respiratory, Thoracic and Mediastinal Disorders:* throat irritation

*Vascular disorders:* hypertension

\*Reported in the 52 week safety study (P04139)

Electrocardiogram QT prolongation occurred at the same incidence as placebo (<1%).

### **Post-Market Adverse Drug Reactions**

No post-marketing safety data is available since ZENHALE™ has not yet been marketed.

## **DRUG INTERACTIONS**

### **Drug-Drug Interactions**

In clinical studies, concurrent administration of ZENHALE™ and other drugs, such as short-acting beta<sub>2</sub>-agonist and intranasal corticosteroids have not resulted in an increased frequency of adverse drug reactions. No formal drug interaction studies have been performed with ZENHALE™. The drug interactions of the combination are expected to reflect those of the individual components.

**Table 2: Established or potential Drug-Drug Interactions**

<b>Drug</b>	<b>Ref.</b>	<b>Effect</b>	<b>Clinical Comment</b>
<i>Ketoconazole</i>	CT	Co-administration of inhaled mometasone furoate with the potent CYP3A4 enzyme inhibitor ketoconazole causes an increase in plasma concentration of mometasone furoate.	Caution should be exercised when considering the coadministration of ZENHALE™ with ketoconazole.
<i>Adrenergic agents</i>	T	Concomitant administration of other sympathomimetic agents may potentiate the undesirable effects of formoterol.	Apart from the recommended use of short-acting beta <sub>2</sub> -agonist rescue medication, caution should be exercised when considering the coadministration of ZENHALE™ with other adrenergic agents.
<i>Xanthine derivatives and diuretics</i>	T	Concomitant treatment with xanthine derivatives or non-potassium sparing diuretics may potentiate the possible hypokalaemic effect of beta <sub>2</sub> -agonists (see WARNINGS AND PRECAUTIONS).	Caution should be exercised when considering the coadministration of ZENHALE™ with xanthine derivatives or non-potassium sparing diuretics.
<i>Monoamine oxidase inhibitors, tricyclic antidepressants and drugs known to prolong the QTc interval</i>	T	Action of adrenergic agonists on the cardiovascular system may be potentiated by these agents.  Drugs that are known to prolong the QTc-interval have an increased risk of ventricular arrhythmia (see WARNINGS AND PRECAUTIONS).	Formoterol, as other beta <sub>2</sub> -agonists, should be administered with caution to patients being treated with these drugs.
<i>Beta-adrenergic receptor antagonists</i>	T	Beta-adrenergic blockers may weaken or antagonize the effect of formoterol.	ZENHALE™ should not be given together with beta-adrenergic blockers (including eye drops) unless there are compelling reasons for their use.
Legend: C=Case Study; CT = Clinical Trial; T = Theoretical			

## DOSAGE AND ADMINISTRATION

### Dosing Considerations

Long-acting beta<sub>2</sub>-agonists (LABA), such as formoterol, one of the active ingredients in ZENHALE™, may increase the risk of asthma-related death. Available data from controlled clinical trials suggest that LABA increase the risk of asthma-related hospitalization in pediatric and adolescent patients (see WARNINGS AND PRECAUTIONS). Therefore, when treating patients with asthma, physicians should only prescribe ZENHALE™ for patients not adequately controlled on a long-term asthma control medication, such as an inhaled corticosteroid or whose disease severity clearly warrants initiation of treatment with both an inhaled corticosteroid and LABA. ZENHALE™ should not be used in patients whose asthma can be managed by occasional

use of short-acting, inhaled beta<sub>2</sub>-agonists or for patients whose asthma can be successfully managed by inhaled corticosteroids along with occasional use of inhaled short-acting beta<sub>2</sub>-agonists. Once asthma control has been achieved and maintained, assess the patient at regular intervals and do not use ZENHALE™ for patients whose asthma can be adequately controlled on low or medium dose inhaled corticosteroids.

Patients should be made aware that for optimum benefit, ZENHALE™ should be taken regularly, even when they are asymptomatic. Rescue inhalations only need to be taken to relieve acute asthma symptoms (see WARNINGS AND PRECAUTIONS).

A reassessment of asthma therapy should be considered in patients using an increasing number of rescue inhalations for symptom relief without achieving improved asthma control.

ZENHALE™ therapy should not be initiated to treat an asthma exacerbation.

Do not use more than two inhalations twice daily of the prescribed strength of ZENHALE™ as some patients are more likely to experience adverse effects with higher doses of formoterol. If symptoms arise between doses, an inhaled short-acting beta<sub>2</sub>-agonist should be taken for immediate relief.

### **Recommended Dose and Dosage Adjustment**

ZENHALE™ is available in three strengths, 50/5, 100/5 and 200/5; delivering 50 mcg, 100 mcg and 200 mcg of mometasone furoate (MF), respectively and 5 mcg of formoterol fumarate dihydrate (F) per actuation.

ZENHALE™ should be administered as two inhalations twice daily (morning and evening) by oral inhalation according to the dose recommended in Table 3.

**Table 3: Recommended dose for ZENHALE™ treatment based on prior asthma therapy**

<b>Previous Therapy</b>	<b>Recommended Dose</b>	<b>Maximum Recommended Daily Dose</b>
Inhaled low dose glucocorticosteroids	ZENHALE™ 50/5, 2 inhalations twice daily	200 / 20 mcg
Inhaled medium dose glucocorticosteroids	ZENHALE™ 100/5, 2 inhalations twice daily	400 / 20 mcg
Inhaled high dose glucocorticosteroids	ZENHALE™ 200/5, 2 inhalations twice daily	800 / 20 mcg

For patients who have not previously received inhaled corticosteroids, but whose disease severity warrants initiation of treatment with two maintenance therapies, depending upon asthma severity, the recommended starting dose must be decided by the physician.

The maximum daily recommended dose is 800 mcg MF / 20 mcg F (given as two inhalations of ZENHALE™ 200/5 twice daily) for patients 12 years of age and older. In the event of an acute attack or if symptoms arise between doses, an inhaled short-acting beta<sub>2</sub>-agonist should be taken for immediate relief. After asthma stability has been achieved, it is desirable to titrate to the lowest effective dosage. Patients should be regularly assessed by a doctor.

If a previously effective dosage regimen of ZENHALE™ fails to provide adequate control of asthma, medical advice should be sought and therapy reassessed, as this may indicate a worsening of the underlying condition.

### **Missed Dose**

If a dose is missed, the patient should be instructed to take the next dose as soon as they remember unless it is near to the time for the next dose, at which time they should wait until the next dose is due. The patient should be instructed not to double the dose.

### **Administration**

ZENHALE™ should be administered by oral inhalation only.

Patients should be made aware that ZENHALE™ must be used daily for optimum benefit, even when asymptomatic.

As a general rule, rinsing the mouth and gargling with water after each inhalation can help in preventing the occurrence of candidiasis. Cleansing dentures has the same effect.

### ***Priming and Use***

Patients should be instructed to properly use the inhaler, as described in PART III: CONSUMER INFORMATION and to read the Product Insert contained in their package.

ZENHALE™ should be administered only by orally inhaled routes. After each dose, the patient should be advised to rinse their mouth out with water without swallowing. The inhaler must be shaken prior to each inhalation.

ZENHALE™ should be primed before using for the first time by removing the mouthpiece and releasing 4 test sprays into the air, away from the face, shaking well before each spray. In cases where the inhaler has not been used for more than 5 days, the inhaler should be primed again as described above.

The ZENHALE™ canister should only be used with the ZENHALE™ actuator. The ZENHALE™ actuator should not be used with any other inhalation drug product. Actuators from other products should not be used with the ZENHALE™ canister.

The correct amount of medication in each inhalation cannot be ensured after the labeled number of actuations from the canister has been used, even though the inhaler may not feel completely empty and may continue to operate. The inhaler should be discarded when the labeled number of actuations has been used (the dose counter will read “0”). Never immerse the canister into water to determine the amount remaining in the canister (“float test”).

### *Use with a spacer device*

Use of the AeroChamber Plus Anti-Static valved holding chamber is recommended with ZENHALE™, in patients who find it difficult to synchronize aerosol actuation with inspiration of breath.

### **OVERDOSAGE**

The risks associated with overdosage for the individual components apply to ZENHALE™.

- Inhalation or oral administration of excessive doses of corticosteroids may lead to suppression of HPA axis function.
- An overdose of formoterol fumarate is likely to lead to effects that are typical of beta<sub>2</sub>-adrenergic stimulants: nausea, vomiting, headache, tremor, drowsiness, palpitations, tachycardia, ventricular arrhythmias, metabolic acidosis, hypokalaemia, hyperglycaemia.

### ***Treatment***

Supportive and symptomatic treatment is indicated. In serious cases, patients should be hospitalised. Use of cardioselective beta-blockers may be considered, but only under the supervision of a physician and with extreme caution since the use of beta-adrenergic blocker medication may provoke bronchospasm. Adrenal function monitoring should be included as part of management.

**For management of suspected drug overdose, contact your regional Poison Control Centre.**

## **ACTION AND CLINICAL PHARMACOLOGY**

### **Mechanism of Action**

ZENHALE™ contains both mometasone furoate and formoterol fumarate; therefore, the mechanisms of actions described below for the individual components apply to ZENHALE™.

### ***Mometasone furoate***

Mometasone furoate is a topical glucocorticosteroid with local anti-inflammatory properties.

Glucocorticoids, like mometasone furoate exert their anti-inflammatory effects through glucocorticoid receptors (GRs). On binding the glucocorticoid, the GR heterocomplex dissociates, and the ligand activated GR translocates from the cytoplasm to the nucleus. The activated GR may then upregulate the transcription of anti-inflammatory genes by binding to specific DNA sequences termed glucocorticoid response elements. However, it is more likely that the primary anti-inflammatory activity of glucocorticoids results from their ability to suppress the transcription of genes. In this case, the activated GR interacts with transcription factors apolipoprotein 1 (AP 1) or nuclear factor kappa B (NF-κB) to down regulate gene expression. In addition, glucocorticoids have been shown to upregulate the expression of an inhibitor of NF-κB.

### ***Formoterol fumarate***

Formoterol fumarate is a potent selective beta<sub>2</sub>-adrenergic stimulant. It exerts a bronchodilator effect in patients with reversible airway obstruction. The effect sets in rapidly and is still significant 12 hours after inhalation. *In vitro*, formoterol inhibits the release of histamine and leukotrienes from passively sensitized human lung. Some anti-inflammatory properties, such as inhibition of oedema and inflammatory cell accumulation, have been observed in animal experiments.

### **Pharmacodynamics**

#### **Mometasone furoate**

Affinity for binding to the GR corresponds to functional activity. Mometasone furoate binds with very high affinity to the human GR which in cells, results in potent inhibition of the synthesis and release of proinflammatory mediators and cytokines.

Mometasone furoate significantly inhibits the release of leukotrienes from leucocytes of allergic patients. In cell culture, mometasone furoate inhibits synthesis and release of IL-1, IL-5, IL-6, and TNF $\alpha$  with high potency; it is also a potent inhibitor of the production of the TH2 cytokines, IL-4 and IL-5, from human CD4<sup>+</sup> T-cells. In mixed leukocytes from atopic patients, mometasone furoate was a more potent inhibitor of leukotriene production than beclomethasone dipropionate.

In preclinical models, mometasone furoate has been shown to reduce the accumulation of inflammatory cells, including eosinophils, infiltrating into the upper and lower airways and improve lung function following allergen provocation. Additionally, mometasone furoate reduces the number of lymphocytes and the levels of messenger RNA for the proallergic cytokines IL-4 and IL-5.

#### **Formoterol fumarate**

*In vitro* studies on guinea pig trachea have indicated that racemic formoterol and its (R,R)- and (S,S)-enantiomers are highly selective beta<sub>2</sub>-adrenoceptor agonists. The (S,S)-enantiomer was 800 to 1,000 times less potent than the (R,R)-enantiomer and did not affect the activity of the (R,R)-enantiomer on tracheal smooth muscle. No pharmacological basis for the use of one of the two enantiomers in preference to the racemic mixture has been demonstrated.

### ***Clinical Safety***

In patients 12 years of age and older with asthma, there was no evidence of significant hypokalemia or hyperglycemia in response to formoterol treatment after doses of formoterol fumarate ranging from 10 microgram to 40 microgram from ZENHALE<sup>™</sup>. No relevant changes in heart rate during the study were observed with ZENHALE<sup>™</sup>. No patients had a QTcB (QTc corrected by Bazett's formula)  $\geq$ 500 msec during treatment. There were no other clinically significant abnormalities or changes in ECG data.

The effects of inhaled mometasone furoate administered via ZENHALE<sup>™</sup> on adrenal function were evaluated in two clinical studies in patients with asthma. HPA-axis function was assessed by 24-hour plasma cortisol AUC. Dose related decreases in plasma cortisol were observed with ZENHALE<sup>™</sup> but these effects are not considered to be clinically significant.

## **Pharmacokinetics**

In a single-dose cross-over study, there was no evidence of a significant pharmacokinetic interaction between mometasone furoate and formoterol when given as ZENHALE™.

### **Absorption:**

**Mometasone furoate:** Following inhalation of single and multiple doses of ZENHALE™, mometasone furoate (200 to 800 mcg) was rapidly absorbed with a prolonged absorption phase. Median  $T_{max}$  values ranged from 0.50 to 4 hours. Exposure to mometasone furoate increased with increasing inhaled dose. Absorbed mometasone furoate is rapidly cleared from plasma at a rate of approximately 12.5 mL/min/kg, independent of dose. The effective  $t_{1/2}$  for mometasone furoate following inhalation with ZENHALE™ was 25 hours. Using the steady-state exposure to mometasone furoate when administered by inhalation from ZENHALE™ and after a single IV dose from different studies, estimates of the absolute bioavailability were approximately 14% in healthy subjects and ranged from 5% to 7% in asthmatic patients.

**Formoterol fumarate:** Following ZENHALE™ administration formoterol was rapidly absorbed with median  $T_{max}$  values ranging from 0.17 to 1.97 hours. Over the dose range of 10 to 40 mcg for formoterol from ZENHALE™, the exposure to formoterol was dose proportional. The mean  $t_{1/2}$  for formoterol in plasma was 9.1 hours.

### **Distribution:**

**Mometasone furoate:** After intravenous bolus administration, the mean steady-state volume of distribution ( $V_d$ ) was 152 liters. The *in vitro* protein binding for mometasone furoate was reported to be 98 to 99% (in a concentration range of 5 to 500 ng/mL).

**Formoterol fumarate:** The plasma protein binding of formoterol was 61% to 64% and binding to human serum albumin was 34%.

### **Metabolism:**

**Mometasone furoate:** Mometasone furoate is extensively metabolized in all species investigated. No major metabolites have been identified. The portion of an inhaled mometasone furoate dose that is swallowed and absorbed from the gastrointestinal tract undergoes extensive metabolism to multiple metabolites. In human liver microsomes, mometasone furoate is metabolized to many metabolites, including 6-beta hydroxy mometasone furoate, which is formed by cytochrome P450 3A4.

**Formoterol fumarate:** Formoterol is eliminated primarily by metabolism, with direct glucuronidation being the major pathway of biotransformation. O-demethylation followed by glucuronidation is another pathway. Minor pathways involve sulphate conjugation of formoterol and deformylation followed by sulphate conjugation. Multiple isozymes catalyse the glucuronidation (UGT1A1, 1A3, 1A6, 1A7, 1A8, 1A9, 1A10, 2B7 and 2B15) and O-demethylation (CYP2D6, 2C19, 2C9 and 2A6) of formoterol, suggesting a low potential for drug-drug interactions through inhibition of a specific isozyme involved in formoterol metabolism. Formoterol did not inhibit cytochrome P450 isozymes at therapeutically relevant concentrations.

**Excretion:**

**Mometasone furoate:** A radiolabeled, orally inhaled dose is excreted mainly in the feces (74%) and to a lesser extent in the urine (8%).

**Formoterol fumarate:** Following oral administration of 80 microgram of radiolabeled formoterol fumarate to 2 healthy subjects, 59% to 62% of the radioactivity was eliminated in the urine and 32% to 34% in the feces over a period of 104 hours. In an oral inhalation study with ZENHALE™, renal clearance of formoterol from the blood was 217 ml/min. Following single inhaled doses of formoterol ranging from 10 to 40 microgram from ZENHALE™, 6.2% to 6.8% of the formoterol dose was excreted in urine unchanged.

**Special Populations and Conditions**

**Pediatrics:** The pharmacokinetics of ZENHALE™ has not been specifically studied in children below 12 years of age.

**Geriatrics:** The pharmacokinetics of ZENHALE™ has not been specifically studied in the elderly population.

**Gender:** Studies to examine the effects of gender on the pharmacokinetics of ZENHALE™ have not been specifically conducted.

Based on analysis of single and multiple dose pharmacokinetics studies, no effect of gender on mometasone furoate and formoterol exposure was observed.

**Race:** Studies to examine the effects of race on the pharmacokinetics of ZENHALE™ have not been specifically conducted.

**Hepatic Insufficiency:** The pharmacokinetics of ZENHALE™ has not been specifically studied in patients with hepatic impairment. Concentrations of mometasone furoate appear to increase with severity of hepatic impairment. These increases are not considered to be clinically significant.

A study evaluating the administration of a single inhaled dose of 400 mcg mometasone furoate by a dry powder inhaler to subjects with mild (n=4), moderate (n=4), and severe (n=4) hepatic impairment resulted in only 1 or 2 subjects in each group having detectable peak plasma concentrations of mometasone furoate (ranging from 50-105 pg/mL). The observed peak plasma concentrations appear to increase with severity of hepatic impairment; however, the numbers of detectable levels were few.

**Renal Insufficiency:** The pharmacokinetics of ZENHALE™ has not been specifically studied in patients with renal impairment.

## **STORAGE AND STABILITY**

Store at controlled room temperature 15°-30°C. Do not freeze. For best results, the inhaler should be warmed to room temperature before use (e.g. by warming in the hands) if it is exposed to low temperatures.

For best results, the canister should be at room temperature before use. Shake well before using. Keep out of reach of children. Avoid spraying in eyes.

Contents under pressure. Do not place in hot water or near radiators, stoves or other sources of heat. Do not puncture or incinerate container or store at temperatures over 50°C.

## **SPECIAL HANDLING INSTRUCTIONS**

No special requirements.

Any unused product or waste should be disposed of in accordance with local requirements.

## **DOSAGE FORMS, COMPOSITION AND PACKAGING**

### **Dosage Forms**

ZENHALE™ is a pressurized metered dosed suspension for inhalation available in three strengths.

- ZENHALE™ 50 /5 delivers 50 microgram of mometasone furoate and 5 microgram of formoterol fumarate dihydrate per actuation.
- ZENHALE™ 100 /5 delivers 100 microgram of mometasone furoate and 5 microgram of formoterol fumarate dihydrate per actuation.
- ZENHALE™ 200 /5 delivers 200 microgram of mometasone furoate and 5 microgram of formoterol fumarate dihydrate per actuation.

### **Composition**

The formulation is a white to off-white suspension. The active ingredients; mometasone furoate and formoterol fumarate dihydrate, are suspended in a mixture of ethanol anhydrous, oleic acid and HFA-227 (propellant).

After priming each actuation the inhaler delivers 60.5, 121.0 or 242.0 microgram of mometasone furoate and 6.1 microgram of formoterol fumarate dihydrate from the valve and delivers 50, 100 or 200 microgram of mometasone furoate and 5 microgram of formoterol fumarate dihydrate from the actuator.

### **Packaging**

The suspension is contained in an aluminium canister internally coated with fluorinated ethylene/propylene copolymer (FEP) and sealed with a 50 microliter metering valve. A polypropylene actuator is provided with the pressurized canister to deliver a dose to the patient. A polypropylene mouthpiece cover is provided with the actuator. The actuator has an integrated dose counter, which shows how many actuations are left. One pressurized container delivers 120 actuations to the patient. There is one inhaler in each package.

## PART II: SCIENTIFIC INFORMATION

### PHARMACEUTICAL INFORMATION


#### Drug Substance

Proper name: ***Mometasone furoate***

Chemical name: 9,21-dichloro-11 $\beta$ ,17-dihydroxy-16 $\alpha$ -methylpregna-1,4-diene-3,20-dione 17-(2)-furoate

Molecular formula  
and molecular mass: C<sub>27</sub>H<sub>30</sub>Cl<sub>2</sub>O<sub>6</sub> 521.44

Structural formula:



Physicochemical  
properties: white powder; practically insoluble in water; slightly soluble in methanol, ethanol, and isopropanol; soluble in acetone.


#### Drug Substance

Proper name: ***Formoterol fumarate dihydrate***

Chemical name: ( $\pm$ )-2-hydroxy-5-[(1RS)-1-hydroxy-2-[[[(1RS)-2-(4-methoxyphenyl)-1-methylethyl]-amino]ethyl]formanilide fumarate dihydrate

Molecular formula  
and molecular mass: (C<sub>19</sub>H<sub>24</sub>N<sub>2</sub>O<sub>4</sub>)<sub>2</sub>•C<sub>4</sub>H<sub>4</sub>O<sub>4</sub>•2H<sub>2</sub>O 840.9

Structural formula:



Physicochemical properties: white to yellowish powder, freely soluble in glacial acetic acid, soluble in methanol, sparingly soluble in ethanol and isopropanol, slightly soluble in water, and practically insoluble in acetone, ethyl acetate, and diethyl ether.

## CLINICAL TRIALS

The safety and efficacy of ZENHALE™ were demonstrated in three randomized, double-blind, parallel group, multicenter clinical studies (P04073, P04334, and P04431) of 12 to 26 weeks in duration involving 2,255 patients 12 years of age and older. Patients with persistent asthma uncontrolled on low, medium or high dose inhaled corticosteroids (baseline FEV1 means of 66% to 75% of predicted normal) were enrolled in ZENHALE™ 50/5, ZENHALE™ 100/5, or ZENHALE™ 200/5 studies, respectively. All of the studies included a 2 to 3-week run-in period with mometasone furoate to establish a level of asthma control consistent with current medical practice. Two clinical trials evaluated ZENHALE™ compared with its individual components, mometasone furoate and formoterol and one clinical study evaluated two different strengths of ZENHALE™ compared with mometasone furoate alone. The details of each study design are presented in Table 4.

## Study demographics and trial design

**Table 4: Summary of patient demographics for ZENHALE™ (MF/F) Safety and Efficacy clinical trials in asthma**

Study #	Trial design <ul style="list-style-type: none"> <li>○ Duration</li> <li>• Primary Endpoints</li> </ul>	Dosage	No. of subjects	Subject Population <ul style="list-style-type: none"> <li>• Age Range (mean)</li> <li>• Male / Female</li> <li>• Mean FEV1 / Mean Percent Predicted FEV1 at Baseline</li> </ul>
P04073	Randomized Multi-center Double-blind Parallel group Placebo controlled <ul style="list-style-type: none"> <li>○ 26 weeks</li> <li>• The change from Baseline to Week 12 in FEV1 AUC(0-12 hr)</li> <li>• Clinically judged asthma deteriorations or reduction in lung function</li> </ul>	MF/F 50/5 MDI: 2 inhalations BID MF 50mcg MDI: 2 inhalations BID F 5 mcg: 2 inhalations BID Placebo: 2 inhalations BID	182  188  188  188  Total = 746	adult and adolescent subjects with persistent asthma previously treated with low doses of ICS <ul style="list-style-type: none"> <li>• 12-79 (38)</li> <li>• 333 / 413</li> <li>• 2.50L / 75%</li> </ul>
P04334	Randomized Multi-center Double-blind Parallel group Placebo controlled <ul style="list-style-type: none"> <li>○ 26 weeks</li> <li>• The change from Baseline to Week 12 in FEV1 AUC(0-12 hr)</li> <li>• Clinically judged asthma deteriorations or reduction in lung function</li> </ul>	MF/F 100/5 MDI: 2 inhalations BID MF 100 mcg MDI: 2 inhalations BID F 5mcg MDI: 2 inhalations BID Placebo: 2 inhalations BID	191  192  202  196  Total = 781	adult and adolescent subjects with persistent asthma previously treated with medium doses of ICS <ul style="list-style-type: none"> <li>• 12-76 (42)</li> <li>• 321 / 460</li> <li>• 2.33L / 73%</li> </ul>
P04431	Randomized Multi-center Double-blind Parallel group Non-placebo controlled <ul style="list-style-type: none"> <li>○ 12 weeks</li> <li>• The change from Baseline to Week 12 in FEV1 AUC(0-12 hr)</li> </ul>	MF/F 100/5 MDI: 2 inhalations BID MF/F 200/5 MDI: 2 inhalations BID MF 200mcg MDI: 2 inhalations BID	233  255  240  Total = 728	adult and adolescent subjects with persistent asthma previously treated with high doses of ICS <ul style="list-style-type: none"> <li>• 12-84 (48)</li> <li>• 319 / 409</li> <li>• 2.05L / 66%</li> </ul>

MF/F = mometasone furoate/formoterol fumarate combination (ZENHALE™); MF = mometasone furoate; F = formoterol fumarate; MDI = metered dose inhaler; BID = twice daily; ICS = inhaled corticosteroid.

## Study results

Superior efficacy was established with ZENHALE™ in all primary endpoints measuring lung function, and asthma deteriorations and in key secondary endpoints measuring nocturnal awakenings and asthma control.

### *Primary Endpoints*

The change from Baseline to Week 12 in FEV1 AUC(0-12 hr) was a co-primary efficacy variable for studies P04073 and P04334 and the primary efficacy variable in study P04431. Clinically judged asthma deteriorations or reduction in lung function was a co-primary endpoint in Studies P04073 and P04334.

In the two 26-week studies, (P04073, P04334) patients who received ZENHALE™ 50/5 or 100/5 had statistically significant greater improvement from baseline in mean FEV1 AUC compared with mometasone furoate (the primary treatment comparison) and versus placebo at Week 12 (both  $p \leq 0.001$ ). This improvement continued through Week 26.

In a 12-week study (P04431) in patients with persistent asthma and prior clinically judged asthma deteriorations or reduction in lung function, patients who received ZENHALE™ 100/5 and ZENHALE™ 200/5 had a statistically significant greater improvement from baseline in mean FEV1 AUC compared to mometasone furoate 200 microgram ( $p < 0.001$ ). Patients receiving ZENHALE™ 200/5 had a greater numerical increase in serial FEV1 from baseline as compared with patients receiving ZENHALE™ 100/5 across the 12 week treatment period.

**Table 5: FEV<sub>1</sub> AUC (0-12 hours) Results of ZENHALE™ (MF/F) Clinical Efficacy Program**  
**[MEAN FEV<sub>1</sub> over 12 hours in Liter<sup>a</sup> units and FEV<sub>1</sub> AUC in Liter (L) x Hour (hr) units]**

Treatment Group	N <sup>b</sup>	Change from Baseline at Week 12			Significant p-values (MF/F vs. treatment arms)
		Mean FEV <sub>1</sub> AUC(0-12 hr) in Liters (% change from baseline) <sup>a</sup>	FEV <sub>1</sub> AUC(0-12 hr) in L x hr		
			Change from baseline	diff. from MF/F	
<b>P04073</b>					
MF/F 50/5 MDI: 2 inhalations BID	155	0.33L (13.8%)	4.00	–	–
MF 50 mcg MDI: 2 inhalations BID	156	0.21L (9.0%)	2.53	1.47	MF/F vs. MF, p=0.001
F 5 mcg MDI: 2 inhalations BID	146	0.32L (12.3%)	3.83	0.17	NS
Placebo: 2 inhalations BID	129	0.09L (4.1%)	1.11	2.89	MF/F vs. Placebo, p<0.001
<b>P04334</b>					
MF/F 100/5 MDI: 2 inhalations BID	166	0.26L (11.7%)	3.11	–	–
MF 100 mcg MDI: 2 inhalations BID	169	0.11L (5.7%)	1.30	1.81	MF/F vs. MF, p<0.001
F 5 mcg MDI: 2 inhalations BID	135	0.16L (8.5%)	1.93	1.18	NS
Placebo: 2 inhalations BID	128	0.05L (3.9%)	0.57	2.54	MF/F vs. Placebo, p<0.001
<b>P04431</b>					
MF/F 100/5 MDI: 2 inhalations BID	204	0.30L (14.4%)	3.59	–	–
MF/F 200/5 MDI: 2 inhalations BID	231	0.35L (16.8%)	4.19	–	–
MF 200 mcg MDI: 2 inhalations BID	211	0.17L (7.6%)	2.04	1.55 2.15	MF/F 100/5 vs. MF, p<0.001 MF/F 200/5 vs. MF, p<0.001

MF/F = mometasone furoate/formoterol fumarate combination (ZENHALE™); MF = mometasone furoate; F = formoterol fumarate; MDI = metered dose inhaler; FEV<sub>1</sub> = forced expiratory volume (Liters) in one second; AUC = area under the curve

<sup>a</sup> Mean FEV<sub>1</sub> over 12 hours in Liters = FEV<sub>1</sub> AUC(0-12 hr) in L x hours divided by the 12-hour serial evaluation interval.

<sup>b</sup> N is the number of subjects with Week 12 data.

In studies P04073 and P04334 patients receiving ZENHALE™ 50/5 or 100/5 had statistically significantly fewer clinically judged asthma deterioration or reduction in lung function compared with patients who received formoterol (the primary treatment comparison) or placebo (p<0.001). Patients who received ZENHALE™ 50/5 or 100/5 also had a delayed time to first clinically

judged asthma deterioration or reduction in lung function compared to patients who received formoterol (the primary treatment comparison) or placebo ( $p < 0.001$ ). The results are detailed in Table 6.

**Table 6: Asthma deteriorations [clinically judged asthma deterioration or reduction in lung function] results**

<b>Study P04073</b>				
<b>Asthma Deteriorations N (%)</b>	<b>MF/F MDI 100/10 µg BID n=182</b>	<b>MF MDI 100 µg BID n=188</b>	<b>F MDI 10 µg BID n=188</b>	<b>Placebo n=188</b>
All First Asthma Deteriorations*	30 (16.5%)	53 (28.2%)	84 (44.7%)	86 (45.7%)
Decrease in FEV1**	6 (3.3%)	14 (7.4%)	28 (14.9%)	30 (16.0%)
Decrease in PEF <sup>++</sup>	23 (12.6%)	36 (19.1%)	46 (24.5%)	43 (22.9%)
Emergency treatment	0	0	2 (1.1%)	1 (0.5%)
Hospitalization	0	0	0	0
Treatment with excluded asthma medication <sup>§</sup>	2 (1.1%)	4 (2.1%)	15 (8.0%)	15 (8.0%)
<b>Study P04334</b>				
<b>Asthma Deteriorations N (%)</b>	<b>MF/F MDI 200/10 µg BID n=192</b>	<b>MF MDI 200 µg BID n=192</b>	<b>F MDI 10 µg BID n=202</b>	<b>Placebo n=196</b>
All First Asthma Deteriorations*	58 (30%)	65 (34%)	109 (54%)	109 (56%)
Decrease in FEV1**	18 (9%)	19 (10%)	31 (15%)	41 (21%)
Decrease in PEF <sup>++</sup>	37 (19%)	41 (21%)	62 (31%)	61 (31%)
Emergency treatment	0	1 (<1%)	4 (2%)	1 (<1%)
Hospitalization	1 (<1%)	0	0	0
Treatment with excluded asthma medication <sup>‡</sup>	2 (1%)	4 (2%)	17 (8%)	8 (4%)

\*Includes only the first event of clinically judged asthma deterioration or reduction in lung function for each patient. Patients could have experienced more than one event criterion on the day of the event.

\*\*Decrease in absolute FEV1 below the treatment period stability limit (defined as 80% of the average of the two predose FEV1 measurements taken 30 minutes and immediately prior to the first dose of randomized trial medication)

<sup>++</sup>Decrease in AM or PM peak expiratory flow (PEF) on 2 or more consecutive days below the treatment period stability limit (defined as 70% of the AM or PM PEF obtained over the last 7 days of the run-in period)

<sup>§</sup>Thirty-Four patients received systemic glucocorticosteroids; 1 patient in F MDI 10 µg BID group and 1 patient in Placebo group received medications other than systemic glucocorticosteroids.

<sup>‡</sup>Thirty patients received systemic glucocorticosteroids; 1 patient in the F MDI 10 µg BID group received medications other than systemic glucocorticosteroids.

### ***Secondary endpoints***

P04073 & P04334: Clinically important improvement in asthma specific quality of life (as measured by Asthma Quality of Life Questionnaire [AQLQ(S) 12+]) and asthma control (as measured by Asthma Control Questionnaire [ACQ]) were observed in patients receiving ZENHALE™ 50/5 or 100/5 as compared with patients receiving placebo. At study endpoint, patients receiving ZENHALE™ 50/5 or 100/5 had a clinically important improvement in asthma specific quality of life compared to placebo with an observed difference of 0.62 point [95% CI 0.44, 0.81] or 0.50 point [95% CI 0.32, 0.68] respectively in AQLQ(S) 12+ score. Patients receiving ZENHALE™ 50/5 or 100/5 were more likely to have well-controlled asthma at study endpoint compared with patients receiving placebo (observed difference was 0.60 point [95% CI 0.43, 0.78] or 0.54 point [95% CI 0.39, 0.70] respectively in ACQ score).

Patients receiving ZENHALE™ 50/5 or 100/5 had statistically significantly greater reduction in proportion of nights with nocturnal awakenings due to asthma that require use of SABA during the 26 weeks treatment duration compared to those on placebo (the primary treatment comparison) and to formoterol fumarate ( $p \leq 0.035$ ).

Patients who received ZENHALE™ 50/5 or 100/5 had a statistically significant improvement in pre-dose morning PEF compared with patients in all other treatment groups at study endpoint ( $p \leq 0.007$  and  $p \leq 0.008$ , respectively).

Patients who received ZENHALE™ 50/5 or 100/5 demonstrated a statistically significant decrease in short-acting beta<sub>2</sub>-agonist rescue use and a decrease in asthma symptom scores versus patients on placebo (the primary treatment comparator), at study endpoint ( $p \leq 0.004$ ).

### ***Tachyphylaxis***

Following the initial dose of ZENHALE™ 50/5, 100/5 and 200/5, FEV<sub>1</sub> relative to Day 1 baseline improved markedly over the first week of treatment and continued to improve over the 12 and 26 weeks of treatment in all three studies. No reduction in the 12-hour bronchodilator effect was observed with either ZENHALE™ 50/5, 100/5 or 200/5 as assessed by FEV<sub>1</sub> following 12 and 26 weeks of therapy.

## **DETAILED PHARMACOLOGY**

### **Animal**

Non-clinical pharmacology studies were conducted for both individual active products. The results of the nonclinical studies did not identify any unique toxicities with the ZENHALE™ combination or any indication of pharmacodynamic interactions. The findings were consistent with the individual components.

### **Human**

Clinical Pharmacokinetic and Pharmacodynamic studies for the mometasone furoate/formoterol metered dose inhaler indicate the data for the monocomponent dry powder inhaler formulations are relevant to the understanding of the performance of the combination product, and thus to its safety and efficacy. ZENHALE™ when administered from the metered dose inhaler (MDI) was safe and well tolerated with no new safety issues or adverse events identified when compared to the monocomponents.

A single-dose, cross over study in healthy subjects (P03658) confirmed that no significant pharmacokinetic interaction occurs between mometasone furoate and formoterol when co-administered via the metered dose inhaler. See ACTION AND CLINICAL PHARMACOLOGY for more detailed information.

**Systemic Safety:** In a single dose, double blind placebo controlled crossover study in 25 patients with asthma, single dose treatment of 10 mcg formoterol fumarate in combination with 100 mcg or 400 mcg of mometasone furoate delivered via ZENHALE™ 50/5 or 200/5 were compared to formoterol fumarate 10 mcg MDI, formoterol fumarate 12 mcg dry powder inhaler (DPI; nominal dose of formoterol fumarate delivered 10 mcg), or placebo. The degree of bronchodilation at 12 hours after dosing with ZENHALE™ was similar to formoterol fumarate delivered alone via MDI or DPI.

ECGs and blood samples for glucose and potassium were obtained prior to dosing and post dose. No downward trend in serum potassium was observed and values were within the normal range and appeared to be similar across all treatments over the 12 hour period. Mean blood glucose appeared similar across all groups for each time point and no changes were of clinical concern. There was no evidence of significant hypokalemia or hyperglycemia in response to formoterol treatment.

No relevant changes in heart rate during the study were observed with ZENHALE™. No patients had a QTcB (QTc corrected by Bazett's formula)  $\geq 500$  msec during treatment. There were no other clinically significant abnormalities or changes in ECG data.

In a single dose crossover study involving 24 healthy subjects, single dose of formoterol fumarate 10, 20, or 40 mcg in combination with 400 mcg of mometasone furoate delivered via ZENHALE™ were evaluated for safety (ECG, blood potassium and glucose changes). ECGs and blood samples for glucose and potassium were obtained at baseline and post dose. Decrease in mean serum potassium was similar across all three treatment groups (approximately 0.3 mmol/L) and values were within the normal range. No relevant trends towards an increase in mean blood glucose values were observed. No relevant changes in heart rate during the study were observed with ZENHALE™. No subjects had a QTcB  $> 500$  msec during treatment.

Five active- and placebo-controlled studies (study duration ranging from 12, 26, and 52 weeks) evaluated 3381 patients 12 years of age and older with asthma. No clinically meaningful changes were observed in potassium and glucose values in patients receiving ZENHALE™. The effects of ZENHALE™ on heart/pulse rate and blood pressure was comparable to that of the individual component mometasone furoate and formoterol fumarate. No changes in vital signs and ECG parameters suggesting a treatment effect were observed and no clinically significant ECG abnormalities were reported in patients receiving ZENHALE™.

**HPA-axis effects:** The effects of inhaled mometasone furoate administered via ZENHALE™ on adrenal function were evaluated in two clinical studies in patients with asthma. HPA-axis function was assessed by 24-hour plasma cortisol AUC. The effects of ZENHALE™ on plasma cortisol are not considered to be clinically significant.

In a 42-day, open-label, placebo and active-controlled study 60 patients with asthma 18 years of age and older were randomized to receive two inhalations twice daily of 1 of the following treatments: ZENHALE™ 100/5, ZENHALE™ 200/5, fluticasone propionate/salmeterol xinafoate 230 mcg/21 mcg, or placebo. At Day 42, the mean change from baseline plasma cortisol AUC(0-24 hr) was 8%, 22% and 34% lower compared to placebo for the ZENHALE™ 100/5 (n=13), ZENHALE™ 200/5 (n=15) and fluticasone propionate/salmeterol xinafoate 230 mcg/21 mcg (n=16) treatment groups, respectively.

In a 52-week safety study, primary analysis of the plasma cortisol 24-hour AUC was performed on 57 patients with asthma who received 2 inhalations twice daily of ZENHALE™ 100/5, ZENHALE™ 200/5, fluticasone propionate/salmeterol 125/25, or fluticasone propionate/salmeterol 250/25. At Week 52, the mean plasma cortisol AUC(0-24 hr) was 2.2%, 29.6%, 16.7%, and 32.2% lower from baseline for the ZENHALE™ 100/5 (n=18), ZENHALE™ 200/5 (n=20), fluticasone propionate/salmeterol 125/25 (n=8), and fluticasone propionate/salmeterol 250/25 (n=11) treatment groups, respectively.

## MICROBIOLOGY

Not applicable.

## TOXICOLOGY

The toxicity observed in animal studies with mometasone furoate and formoterol fumarate, given in combination as ZENHALE™ or separately, were effects associated with exaggerated pharmacological activity.

### Toxicity Studies

Species	Route of Administration	ZENHALE™ Dose Ratio	Duration
Rats	Inhalation	50:5 and 200:5	2 weeks and 13 weeks
Dogs	Inhalation	50:5 and 200:5	2 weeks and 13 weeks

In 2- and 13-week inhalation toxicity studies conducted in rats and dogs using formulations containing ratios of 50:5 and 200:5 mometasone furoate:formoterol fumarate dihydrate, all findings were consistent with toxicities that would be expected with the individual active drugs. No new or additive toxicities were observed. No pharmacokinetic interactions were observed after co-administration of mometasone furoate and formoterol fumarate.

ZENHALE™ contains both mometasone furoate and formoterol fumarate; therefore, the mutagenicity, carcinogenicity and reproductive toxicity information of the individual components described below apply to ZENHALE™.

Specific mutagenicity, carcinogenicity, and reproduction toxicity studies have not been conducted with ZENHALE™.

## Mutagenicity

### ***Mometasone furoate***

Mometasone furoate was non-mutagenic in the mouse-lymphoma assay and the Salmonella/*E. coli*/mammalian microsome mutagenicity bioassay. At cytotoxic doses only, mometasone furoate produced an increase in chromosome aberrations *in vitro* in Chinese hamster ovary cell (CHO) cultures in the non-activation phase, but not in the presence of rat liver S9 fraction. However, mometasone furoate did not induce chromosomal aberrations *in vitro* in a Chinese hamster lung cell (CHL) chromosomal-aberrations assay, or *in vivo* in the mouse bone marrow erythrocyte-micronucleus assay, in the rat bone-marrow clastogenicity assay, and the mouse male germ-cell clastogenicity assay. Mometasone furoate also did not induce unscheduled DNA synthesis *in vivo* in rat hepatocytes. The finding of simple chromosomal aberrations in the non-activation phase of the CHO assay is considered to be related to cytotoxicity and is not considered to be of significance in the risk assessment of mometasone furoate because of the negative results in the S9 phase of this assay, the negative results in a second *in vitro* chromosomal aberrations assay (CHL assay), and the negative results in three *in vivo* chromosomal aberrations assays.

### ***Formoterol fumarate***

Mutagenicity tests covering a broad range of experimental endpoints have been conducted. No genotoxic effects were found in any of the *in vitro* or *in vivo* tests performed.

## Carcinogenicity

### ***Mometasone furoate***

In a 2-year carcinogenicity study in Sprague Dawley rats, mometasone furoate demonstrated no statistically significant increase in the incidence of tumors at inhalation doses up to 67 mcg/kg (approximately 8 times the maximum recommended daily inhalation dose in adults on an AUC basis and 2 times the maximum recommended daily inhalation dose in pediatric patients based on a mcg/m<sup>2</sup> bases). In a 19-month carcinogenicity study in Swiss CD-1 mice, mometasone furoate demonstrated no statistically significant increase in the incidence of tumors at inhalation doses up to 160 mcg/kg (approximately 10 times the maximum recommended daily inhalation dose in adults on an AUC basis and 2 times the maximum recommended daily inhalation dose in pediatric patients base on a mcg/m<sup>2</sup> bases).

### ***Formoterol fumarate***

Two-year studies in rats and mice did not show any carcinogenic potential.

Male mice treated at very high dose levels showed a slightly higher incidence of benign adrenal subcapsular cell tumours. However, this finding was not seen in a second mouse feeding study, in which pathological changes at high doses consisted of an increased incidence both of benign smooth muscle tumours in the female genital tract, and of liver tumours in both sexes. Smooth muscle tumours are a known effect of beta-agonists given at high doses in rodents.

Two studies in rats, covering different dose ranges, showed an increase in mesovarial leiomyomas. These benign neoplasms are typically associated with long-term treatment of rats at high doses of beta<sub>2</sub>-adrenergic drugs. Increased incidences of ovarian cysts and benign granulosa/theca cell tumours were also seen; beta-agonists are known to have effects on the

ovary in rats which are very likely specific to rodents. A few other tumour types noted in the first study using the higher doses were within the incidences of the historical control population, and were not seen in the lower-dose experiment.

None of the tumour incidences were increased to a statistically significant extent at the lowest dose of the second rat study, a dose leading to a systemic exposure 10 times higher than that expected from the maximum recommended dose of formoterol in humans.

On the basis of these findings and the absence of a mutagenic potential, it is concluded that use of formoterol at therapeutic doses does not present a carcinogenic risk.

### Reproduction toxicity

#### ***Mometasone furoate***

In studies of reproductive function, subcutaneous mometasone furoate was well tolerated at doses up to 7.5 microgram/kg. At 15 microgram/kg, mometasone furoate caused prolonged gestation and prolonged and difficult labor occurred with a reduction in offspring survival and body weight or body weight gain. There was no effect on fertility.

#### ***Formoterol fumarate***

Reproduction studies in rats revealed no impairment of fertility at oral doses up to 3 mg/kg (approximately 1000 times the maximum recommended daily inhalation dose in humans on a mg/m<sup>2</sup> basis).

### Teratogenicity

Like other glucocorticoids, mometasone furoate is a teratogen in rodents and rabbits. Teratology studies were conducted in rats, mice and rabbits by the oral, topical, and/or subcutaneous routes. Effects noted were umbilical hernia in rats, cleft palate in mice, and gall bladder agenesis, umbilical hernia, and flexed front paws in rabbits. There were also reductions in maternal body weight gains, effects on fetal growth (lower fetal body weight and/or delayed ossification) in rats, rabbits and mice, and reduced offspring survival in mice.

In an oral teratology study in rabbits, at 700 µg/kg, increased incidences of resorption and malformation, including cleft palate and/or malformation (hydrocephaly or domed head) were observed. Pregnancy failure was observed in most rabbits at 2800 µg/kg.

## REFERENCES

1. PRODUCT MONOGRAPH: <sup>PR</sup>**FORADIL\*** (formoterol fumarate Dry Powder Capsules for Inhalation)
2. NDS: **AZMAX Twisthaler\*** (Mometasone Furoate Dry Powder Inhaler): Product Monograph dated February 3, 2001
3. Meltzer E et al. Low-dose Mometasone Furoate/Formoterol Therapy: Efficacy and Safety Findings from a Study Investigating a New Combination Therapy in Subjects Whose Asthma was Inadequately Controlled Using Low-Dose Inhaled Corticosteroid. *J Allergy Clin Immunol.* 2010;125(2):AB196.
4. Nathan R et al. Twenty-Six-Week Efficacy and Safety Study of Mometasone Furoate/Formoterol 200/10 µg Combination Treatment in Persistent Asthmatics Previously Uncontrolled on Medium-Dose Inhaled Corticosteroids. *Allergy Asthma Proceedings.* 2010;31(4):269-279.
5. Weinstein S et al. Twelve-Week Efficacy and Safety Study of Mometasone Furoate/Formoterol 200/10 mcg and 400/10 mcg Combination Treatments in Patients With Persistent Asthma Previously Receiving High-Dose Inhaled Corticosteroids. *Allergy Asthma Proc.* 2010;31(4):280-289.
6. Maspero J, et al. Long-Term Safety of Mometasone Furoate/Formoterol Combination for Treatment of Patients With Persistent Asthma. *Journal of Asthma.* 2010;47(10):1106-1115.
7. Bernstein D et al. Non-Inferiority Efficacy Comparison of Mometasone Furoate/Formoterol Versus Fluticasone Propionate/Salmeterol Combination Therapies in Subjects With Persistent Asthma. *Allergy, Asthma & Clinical Immunology.* 2010;6(Suppl 2)

\*All other trademarks are the property of their respective owner(s).

**PART III: CONSUMER INFORMATION****PrZENHALE™**

mometasone furoate / formoterol fumarate dihydrate  
inhalation aerosol

This leaflet is part III of a three-part "Product Monograph" published when ZENHALE™ was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about ZENHALE™. Contact your doctor or pharmacist if you have any questions about the drug.

**ABOUT THIS MEDICATION****What the medication is used for:**

Your doctor has prescribed ZENHALE™ to help control asthma.

ZENHALE™ is used twice a day to help people 12 years of age or older who need treatment for asthma and the doctor recommends a combination product. ZENHALE™ is not a rescue medication. Your doctor should prescribe a rescue medication for you. Use the rescue medication if you have sudden asthma symptoms.

ZENHALE™ is for patients with asthma that:

- was not well controlled with other asthma controlling medications or
- is severe enough that the doctor recommends starting asthma treatment with two asthma medications

This medicine is for **you**. Only a doctor can prescribe it for you. Never give it to someone else. It may harm them, even if their symptoms are the same as yours.

**What it does:**

ZENHALE™ contains two medicines, mometasone furoate and formoterol fumarate dihydrate:

- Mometasone furoate is a corticosteroid. Corticosteroids are used to prevent asthma attacks because they have an anti-inflammatory effect (reduce swelling and irritation in the walls of the small air passages of the lungs, easing breathing problems).
- Formoterol fumarate dihydrate is a long-acting bronchodilator. Bronchodilators help the airways in the lungs to stay open. It makes breathing easier by relaxing muscle spasms in the air passages of the lungs. The effects last for 12 hours.

**When it should not be used:****Do not take ZENHALE™**

- if you are allergic (hypersensitive) to mometasone furoate, formoterol fumarate dihydrate or any of the ingredients contained in the product.
- if you are under 12 years of age.

**What the medicinal ingredients are:**

mometasone furoate and formoterol fumarate dihydrate.

**What the nonmedicinal ingredients are:**

ZENHALE™ is suspended in a propellant (HFA-227), with ethanol and oleic acid.

**What dosage forms it comes in:**

ZENHALE™ is supplied to you in an inhaler containing 50, 100 or 200 micrograms of mometasone furoate and 5 micrograms of formoterol fumarate per actuation (puff).

**WARNINGS AND PRECAUTIONS****Serious Warnings for Asthma Patients**

Formoterol, one of the medicines in ZENHALE™, may increase the risk of asthma-related death. It may increase the risk of asthma-related hospitalization in paediatrics and adolescent patients. Therefore:

- Your doctor will assess your asthma control at regular intervals. ZENHALE™ should only be used when your doctor decides that other asthma medications (e.g. inhaled glucocorticosteroids along with an as needed relief medication) are not helping you enough, or that you need two maintenance medications to control your asthma.
- ZENHALE™ should not be the first medication you use, unless advised by your doctor.

Once asthma control has been achieved and maintained, your doctor will assess you at regular intervals. ZENHALE™ should only be used when an inhaled corticosteroid does not adequately control your asthma symptoms.

BEFORE you use ZENHALE™ talk to your doctor or pharmacist if:

- you are diabetic
- you have high blood pressure or you have ever had a heart problem
- you have liver disease or cirrhosis
- you have problems with your thyroid or adrenal glands
- you have low levels of potassium in your blood
- you have or have ever had tuberculosis
- you have herpes simplex (virus) infection of the eye or any other type of untreated infection

If you are pregnant, planning to get pregnant or are breastfeeding talk to your doctor before using ZENHALE™.

**Important things to remember when using ZENHALE™**

- If symptoms get worse (increased use of a 'reliever' medication peak flow measurement falls, your symptoms are waking you up at night) or do not get better after using ZENHALE™, you should contact your doctor as soon as possible.
- ZENHALE™ should not be used to treat sudden asthma symptoms.

- Do not stop taking ZENHALE™ on your own, even if you feel better. Your doctor can direct you on how to discontinue ZENHALE™ slowly to avoid asthma flare-ups.
- If you are transferring to ZENHALE™ from steroid tablets or syrup and your doctor is slowly decreasing the dose, you may notice that you develop symptoms of allergies such as itchy, watery eyes or rash, which had been controlled by your steroid. Your doctor will tell you how to control these symptoms. During this time, if you begin to notice joint or muscle pain, feelings of depression, tiredness or lethargy, you should tell your doctor.
- You may need to also take steroid tablets or syrup during a severe asthma attack, during other illnesses or during times of stress. Your doctor may give you some steroid tablets or a syrup to carry with you as well as a steroid warning card, which will give you advice on when and how to use them.
- Adolescents 12 years and older treated with ZENHALE™ may grow more slowly than others.
- While using inhaled corticosteroids, patients should avoid coming into contact with anyone who has measles or chicken pox. If you or your child are exposed, tell your doctor right away.
- While using inhaled corticosteroids, patients should be monitored for signs of cataracts, glaucoma (see your eye doctor regularly) or of osteoporosis (decreased bone density).

## INTERACTIONS WITH THIS MEDICATION

Tell your doctor or pharmacist if you are taking or have recently taken any other medications.

### Drugs that may interact with ZENHALE™ include:

- Beta-blocker medicines (such as atenolol or propranolol used for high blood pressure) including eye drops (such as timolol eyedrops for glaucoma)
- Ketoconazole (an antifungal medicine)
- Corticosteroids (by mouth or by injection)
- Diuretics (water pills)
- Xanthine medicines (such as theophylline and aminophylline) used to treat asthma
- Medicines for a fast or uneven heartbeat (such as quinidine)
- Tricyclic anti-depressants
- Monoamine oxidase inhibitors (MAOIs)
- Terfenadine or astemizole (an antihistamine used to treat allergies)
- Other medicines containing a long-acting beta<sub>2</sub>-agonist (i.e. formoterol, salmeterol)
- disopyramide, procainamide or phenothiazines

Make sure that your doctor or pharmacist knows what other medications you are taking, including those you can buy without a prescription and as natural health products.

## PROPER USE OF THIS MEDICATION

Always take ZENHALE™ exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.

As prescribed by your doctor, you should always carry a ‘reliever’ medication such as salbutamol with you. Use your ‘reliever’ medication if your asthma symptoms occur between doses. If your ‘reliever’ medication becomes less effective, seek medical attention right away.

### Usual dose:

Your doctor will decide the best dose for your individual case. The usual dose is 2 puffs in the morning and 2 puffs in the evening. It is important that you use ZENHALE™ every day, even if you have no asthma symptoms at the time.

Do not exceed the dosage of ZENHALE™ recommended to you by your doctor.

### Instructions for use

#### The parts of your ZENHALE™ inhaler

There are two main parts (see figure 1):

- The metal canister that holds the medicine.
- The blue plastic actuator that sprays the medicine from the canister. The mouthpiece of the actuator is covered by a cap.




Figure 1

The inhaler contains 120 actuations (puffs).


The inhaler comes with a dose counter located on the plastic actuator. The dose counter display will show the number of actuations (puffs) of medication remaining. The dose counter will initially display “124” actuations remaining. Each time you press the canister, a puff of medicine is released and the counter will count down by one. The counter will stop counting at 0.

It is recommended not to remove the canister from the actuator because reinsertion may cause the counter to count down by one and/or discharge a puff.

Use the ZENHALE™ canister only with the actuator supplied with the product. Parts of the ZENHALE™ inhaler should not be used with parts from any other inhalation drug product.

**Before using your ZENHALE™ inhaler**

Remove the cap from the mouthpiece of the actuator (see figure 2). Check the mouthpiece for foreign objects and make sure the canister is fully inserted into the actuator.



**Figure 2**


It is necessary to prime the inhaler into the air the first time it is used. Prime the inhaler by releasing four actuations (puffs) into the air, away from your face. Shake the inhaler well before each of the priming actuations. After priming four times, the dose counter should read “120”. It is also necessary to prime the inhaler when it has not been used for more than five days using four actuations (puffs).

**How to use your ZENHALE™ inhaler properly**

Be in an upright position when using ZENHALE™.

**Avoid spraying in your eyes**

1. Remove the cap from the mouthpiece of the actuator.
2. Hold the aerosol can upright between your thumb and forefinger **and shake the inhaler**.
3. Breathe out fully through the mouth as far as is comfortable, expelling as much air from your lungs as possible. Hold the inhaler in the upright position and place the mouthpiece into your mouth. Close your lips around the mouthpiece (see figure 3).



**Figure 3**

4. While breathing in deeply and slowly through your mouth, press down firmly and fully on the canister top until it stops moving in the actuator. Take your finger off the canister.
5. When you have finished breathing in, hold your breath for up to 10 seconds, or for as long as is comfortable. Then remove the inhaler from your mouth and breathe out through your nose while keeping your lips closed.
6. For administration of the second actuation (puff), wait about **30 seconds, shake the inhaler well again and repeat** steps 3 through 5.
7. Firmly replace the cap immediately after use. Do not use excessive force.
8. After completing both inhalations, rinse out your mouth thoroughly with water. Do not swallow the water after rinsing.

**When to replace your ZENHALE™ inhaler**

When the counter reads 20, you should refill your prescription or ask your doctor if you need a new prescription for ZENHALE™.

Discard ZENHALE™ after the counter reaches 0, indicating that you have used the number of actuations on the product label and box. Your inhaler may not feel empty and it may continue to operate, but you will not get the right amount of medicine if you keep using it.

Never try to change the numbers on the counter or remove the counter from the actuator.

**How to clean your ZENHALE™ inhaler**

The mouthpiece should be cleaned using a dry wipe after every 7 days of use.

Routine cleaning instructions:

- Remove the cap off the mouthpiece. Wipe the inside and outside surfaces of the actuator mouthpiece with a clean, dry lint-free tissue or cloth. Put the cap back on the mouthpiece after cleaning.
- Do not attempt to unblock the actuator with a sharp object, such as a pin.
- **Do not wash or put any parts of your inhaler in water.**

**Use with a spacer**

If necessary, your doctor will prescribe the use of a spacer to help you properly take this medication. Instructions provided with the spacer device should be consulted prior to use.

**Overdose:**

If you use more ZENHALE™ than you should, contact your doctor or pharmacist.

The most common symptoms that may occur if you use more ZENHALE™ than you should are nausea, vomiting, headache, trembling, drowsiness, low potassium levels, high sugar levels, or a rapid or irregular heart beat.

If you have used larger doses for a long period of time, you should talk to your doctor or pharmacist for advice. This is because larger doses of ZENHALE™ may reduce the amount of steroid hormones produced by the adrenal gland.

**In case of drug overdose, contact a healthcare practitioner, hospital emergency department or regional poison control centre, even if there are no symptoms.**

**Missed Dose:**

If you miss a dose, take it as soon as you remember. However, if it is nearly time for your next dose, skip the missed dose. Do not take a double dose to make up for the forgotten dose.

**Do not suddenly stop using this medicine even if your asthma seems to be better. Speak with your doctor first.**

Your symptoms may come back if you stop using this medicine before your doctor has told you to do so. If you think that your asthma is not getting any better or seems to be getting worse, after you have started using ZENHALE™ go back and see your doctor.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

**SIDE EFFECTS AND WHAT TO DO ABOUT THEM**

Like all medicines, ZENHALE™ can cause side effects, although not everybody gets them.

Side effects may include nausea, headache, hoarseness, tremor, shakiness, nervousness, dizziness, disturbed sleep, muscle cramps, sore throat, rounded face, loss of bone density and dry mouth.

<b>SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM</b>				
<b>Symptom / effect</b>		<b>Talk with your doctor or pharmacist</b>		<b>Stop taking drug and seek immediate emergency medical attention</b>
		<b>Only if severe</b>	<b>In all cases</b>	
<b>Common</b>	Thrush: a fungal infection in the mouth  Thrush is less likely to occur if you rinse your mouth with water and spit after using ZENHALE™.		√	
<b>Un-common</b>	Fast or uneven heartbeat		√	
	High blood pressure		√	
<b>Rare</b>	Allergic reaction: rash, itching, hives, redness or swelling of the eyes, lips and throat, trouble breathing or joint pain			√
	Increased pressure in the eye (glaucoma) or cataracts (clouding of the lens in the eye), blurry vision, eye pain		√	
	Sudden coughing, wheezing and difficulty breathing (bronchospasm) when you inhale ZENHALE™			√  If you notice this, use your “reliever” inhaler and contact your doctor straightaway.

*This is not a complete list of side effects. For any unexpected effects while taking ZENHALE™, contact your doctor or pharmacist.*

**HOW TO STORE IT**

**Keep your medicine in a safe place out of the reach and sight of children. Your medicine may harm them.**

- Store ZENHALE™ between 15°C to 30°C. If the inhaler is exposed to low temperatures, warm it to room temperature (by warming in the hand) before use.
- Contents under pressure. Do not place in hot water or near radiators, stoves or other sources of heat. Do not puncture or incinerate container or store at temperatures over 50°C.

**WARNING:** Do not use ZENHALE™ after the expiry date which is stated on the carton and the can. The expiry date refers to the last day of that month.

Do not use ZENHALE™ if you notice that the pack is damaged or shows signs of tampering.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

**REPORTING SUSPECTED SIDE EFFECTS**

You can report any suspected adverse reactions associated with the use of health products to the Canada Vigilance Program by one of the following 3 ways:

- Report online at [www.healthcanada.gc.ca/medeffect](http://www.healthcanada.gc.ca/medeffect)
- Call toll-free at 1-866-234-2345
- Complete a Canada Vigilance Reporting Form and:
  - Mail to: Canada Vigilance Program  
Health Canada  
Postal Locator 0701C  
Ottawa, ON K1A 0K9

Postage paid labels, Canada Vigilance Reporting Form and the adverse reaction reporting guidelines are available on the MedEffect™ Canada Web site at [www.healthcanada.gc.ca/medeffect](http://www.healthcanada.gc.ca/medeffect).

*NOTE: Should you require information related to the management of side effects, contact your health professional. The Canada Vigilance Program does not provide medical advice.*

**MORE INFORMATION**

You may need to read this package insert again. **Please do not throw it away** until you have finished your medicine.

This document plus the full product monograph, prepared for health professionals can be found by contacting the sponsor at:

**Merck Canada Inc.**

Kirkland, Quebec H9H 4M7  
1-800-463-5442

This leaflet was prepared by Merck Canada Inc.

Last revised: June 30, 2011

©2011 Merck Canada Inc., a subsidiary of **Merck & Co., Inc.**  
All rights reserved.

™Trademark of Schering-Plough Ltd. Used under license.

