

PRODUCT MONOGRAPH
INCLUDING PATIENT MEDICATION INFORMATION

 **ZERBAXA[®]**

Ceftolozane and Tazobactam powder for injection

1.5 gram (g) per vial

Containing ceftolozane 1 g (as ceftolozane sulfate) and tazobactam 0.5 g (as tazobactam sodium)

Cephalosporin and Beta-lactamase Inhibitor
antibacterial for systemic use

Merck Canada Inc.
16750 route Transcanadienne
Kirkland, QC
Canada H9H 4M7
www.merck.ca

Date of Preparation:
July 26, 2017

Submission Control No: 205115

Table of Contents

PART I: HEALTH PROFESSIONAL INFORMATION.....	3
SUMMARY PRODUCT INFORMATION	3
INDICATIONS AND CLINICAL USE.....	3
CONTRAINDICATIONS	4
WARNINGS AND PRECAUTIONS.....	4
ADVERSE REACTIONS.....	6
DRUG INTERACTIONS	8
DOSAGE AND ADMINISTRATION.....	9
OVERDOSAGE	12
ACTION AND CLINICAL PHARMACOLOGY	13
STORAGE AND STABILITY.....	17
DOSAGE FORMS, COMPOSITION AND PACKAGING	17
PART II: SCIENTIFIC INFORMATION	18
PHARMACEUTICAL INFORMATION.....	18
CLINICAL TRIALS.....	19
MICROBIOLOGY	22
TOXICOLOGY	27
REFERENCES	30
PART III: PATIENT MEDICATION INFORMATION	31



ceftolozane and tazobactam

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form / Strength	Clinically Relevant Nonmedicinal Ingredients
Intravenous Injection	Lyophilized Powder 1.5 gram (g) per vial (ceftolozane 1 g as ceftolozane sulfate and tazobactam 0.5 g as tazobactam sodium)	Sodium (230 mg per vial) <i>For a complete listing see Dosage Forms, Composition and Packaging section.</i>

INDICATIONS AND CLINICAL USE

ZERBAXA[®] (ceftolozane and tazobactam) is indicated for the treatment of patients 18 years of age or older with the following infections when caused by ZERBAXA[®] susceptible strains of the designated microorganisms:

- **Complicated Intra-abdominal Infections**

ZERBAXA[®] is indicated for the treatment of complicated intra-abdominal infections (cIAI) caused by the following Gram-negative and Gram-positive microorganisms: *Enterobacter cloacae*, *Escherichia coli*, *Klebsiella oxytoca*, *Klebsiella pneumoniae*, *Proteus mirabilis*, *Pseudomonas aeruginosa*, *Bacteroides fragilis*, *Streptococcus anginosus*, *Streptococcus constellatus*, and *Streptococcus salivarius*.

Note: In the treatment of cIAI, ZERBAXA[®] should be used in combination with metronidazole in order to provide adequate anaerobic coverage.

- **Complicated Urinary Tract Infections, including Pyelonephritis**

ZERBAXA[®] is indicated for the treatment of complicated urinary tract infections (cUTI), including pyelonephritis caused by the following Gram-negative microorganisms: *Escherichia coli*, *Klebsiella pneumoniae*, *Proteus mirabilis*, and *Pseudomonas aeruginosa*.

To reduce the development of drug-resistant bacteria and maintain the effectiveness of ZERBAXA[®] and other antibacterial drugs, ZERBAXA[®] should be used only to treat infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying

antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

Geriatrics (≥ 65 years of age): Of the 1015 patients treated with ZERBAXA[®] in the Phase 3 clinical trials, 250 (24.6%) were ≥ 65 years of age, including 113 (11.1%) ≥ 75 years. A higher incidence of adverse reactions was observed in patients aged 65 years and older. In the cIAI trial, cure rates were lower in elderly patients. This finding in the elderly population was not observed in the cUTI trial (see **WARNINGS AND PRECAUTIONS** and **ACTION AND CLINICAL PHARMACOLOGY**).

Pediatrics (<18 years of age): Safety and effectiveness in pediatric patients have not been established.

CONTRAINDICATIONS

ZERBAXA[®] is contraindicated in:

- Patients who are hypersensitive to this drug or to any ingredient in the formulation, including any non-medicinal ingredient, or component of the container. For a complete listing, see the **DOSAGE FORMS, COMPOSITION AND PACKAGING** section of the product monograph.
- Patients with known serious hypersensitivity to penicillins, cephalosporins, or other beta-lactams.

WARNINGS AND PRECAUTIONS

Serious Warnings and Precautions

- Serious and occasionally fatal hypersensitivity (anaphylactic) reactions have been reported in patients receiving beta-lactam antibacterials. Before therapy with ZERBAXA[®] is instituted, careful inquiry about previous hypersensitivity reactions to other cephalosporins, penicillins, or beta-lactams should be made. If this product is to be given to a patient with a cephalosporin, penicillin, or other beta-lactam allergy, caution should be exercised because cross sensitivity has been established. If an anaphylactic reaction to ZERBAXA[®] occurs, the drug should be discontinued and appropriate therapy instituted.

Carcinogenesis and Mutagenesis

Long-term carcinogenicity studies in animals have not been conducted with ceftolozane and tazobactam, ceftolozane, or tazobactam (see **TOXICOLOGY**).

Gastrointestinal

***Clostridium difficile*-associated Disease**

Clostridium difficile-associated disease (CDAD) has been reported with the use of nearly all systemic antibacterial agents, including ZERBAXA[®], and may range in severity from mild diarrhea to fatal colitis (see **ADVERSE REACTIONS**). Treatment with antibacterial agents alters the normal flora of the colon which may lead to overgrowth of *C. difficile*.

C. difficile produces toxins A and B which contribute to the development of CDAD. Hypertoxin producing strains of *C. difficile* cause increased morbidity and mortality, as these infections can be refractory to antimicrobial therapy and may require colectomy. CDAD must be considered in all patients who present with diarrhea following antibiotic use. Careful medical history is necessary because CDAD has been reported to occur more than 2 months after the administration of antibacterial agents.

If CDAD is suspected or confirmed, ongoing antibiotic use not directed against *C. difficile* may need to be discontinued. Appropriate fluid and electrolyte management, protein supplementation, antibiotic treatment of *C. difficile*, and surgical evaluation should be instituted as clinically indicated.

Susceptibility/Resistance

Development of Drug Resistant Bacteria

Prescribing ZERBAXA[®] in the absence of a proven or strongly suspected bacterial infection is unlikely to provide benefit to the patient and risks the development of drug-resistant bacteria.

Special Populations

Pregnant Women: There are no adequate and well-controlled trials in pregnant women with either ceftolozane or tazobactam. Tazobactam crosses the placenta in humans. It is not known if ceftolozane crosses the placenta in humans or animals, but it is expected to occur based on the known placental transfer of other cephalosporins. Because animal reproduction studies are not always predictive of human response, ZERBAXA[®] should be used during pregnancy only if the potential benefit to the mother outweighs the possible risk to the fetus.

In animal studies ceftolozane was associated with impaired auditory startle response in rat pups; and tazobactam was associated with maternal toxicity, increased stillbirths and decreased rat pup bodyweight. The reproductive and developmental toxicity of the combination of ceftolozane and tazobactam has not been studied.

In a pre-postnatal study in rats, intravenous ceftolozane administered maternally during pregnancy and lactation (Gestation Day 6 through Lactation Day 20) was associated with a decrease in auditory startle response in postnatal Day 60 male pups at maternal doses of greater than or equal to 300 mg/kg/day. The maternal plasma exposure (AUC) associated with the NOAEL dose of 100 mg/kg/day in rats is approximately 0.4 fold of the mean daily human ceftolozane exposure in healthy adults at the clinical dose of 1 gram thrice-daily.

In a pre-postnatal study in rats, tazobactam administered intraperitoneally twice daily at the end of gestation and during lactation (Gestation Day 17 through Lactation Day 21) produced decreased maternal food consumption and body weight gain at the end of gestation and significantly more stillbirths with a tazobactam dose of 1280 mg/kg/day (approximately 8 times the recommended human dose based on body surface area comparison). No effects on the development, function, learning or fertility of F1 pups were noted, but postnatal body weights for F1 pups delivered to dams receiving 320 and 1280 mg/kg/day tazobactam were significantly reduced 21 days after delivery. F2-generation fetuses were normal for all doses of tazobactam. The NOAEL for reduced F1 body weights was considered to be 40 mg/kg/day (approximately 0.3 times the recommended human dose based on body surface area comparison) (see **TOXICOLOGY**).

Nursing Women: It is not known whether ceftolozane or tazobactam is excreted in human milk. Ceftolozane is expected to be excreted in human milk, based on the known human milk excretion of other cephalosporins. Because many drugs are excreted in human milk, ZERBAXA[®] should not be administered to a nursing woman unless the expected benefit to the mother outweighs the potential risk to the infant (see Pregnant Women; see **TOXICOLOGY**).

Pediatrics (< 18 years of age): Safety and effectiveness in pediatric patients have not been established.

Geriatrics (≥ 65 years of age): Of the 1015 patients treated with ZERBAXA[®] in the Phase 3 clinical trials, 250 (24.6%) were ≥ 65 years of age, including 113 (11.1%) ≥ 75 years. The incidence of adverse events in both treatment groups was higher in older subjects (65 years or older) in the trials for both cUTI and cIAI. In the cIAI trial, cure rates were lower in elderly patients. This finding in the elderly population was not observed in the cUTI trial.

ZERBAXA[®] is substantially excreted by the kidney and the risk of adverse reactions to ZERBAXA[®] may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection in this age group and it may be useful to monitor renal function. Dosage adjustment for elderly patients should be based on renal function (see also **DOSAGE AND ADMINISTRATION** and **ACTION AND CLINICAL PHARMACOLOGY**, Special Populations and Conditions, Geriatrics).

Patients with Renal Impairment: Dosage adjustment is required in patients with moderate (CrCL 30 to 50 mL/min) or severe (CrCL 15 to 29 mL/min) renal impairment and in patients with end-stage renal disease on hemodialysis (see also **DOSAGE AND ADMINISTRATION** and **ACTION AND CLINICAL PHARMACOLOGY**).

ADVERSE REACTIONS

Adverse Reaction Overview

The most common adverse drug reactions (≥ 5% in either indication) occurring in patients receiving ZERBAXA[®] (ceftolozane and tazobactam) in clinical trials were nausea, diarrhea,

headache and pyrexia. The majority of adverse drug reactions were reported as mild to moderate in severity. Treatment discontinuation due to adverse events occurred in 2.0% (20/1015) of patients receiving ZERBAXA[®] and 1.9% (20/1032) of patients receiving comparator drugs. Renal impairment (including the terms renal impairment, renal failure, and renal failure acute) led to discontinuation of treatment in 5/1015 (0.5%) of subjects receiving ZERBAXA[®] and none in the comparator arms.

The overall incidence of mild, moderate, and severe adverse events was similar for the ZERBAXA[®] and comparator groups.

Clinical Trial Adverse Drug Reactions

Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.

ZERBAXA[®] was evaluated in Phase 3 comparator-controlled clinical trials of complicated intra-abdominal infections and complicated urinary tract infections, which included a total of 1015 patients treated with ZERBAXA[®] and 1032 patients treated with comparator (levofloxacin or meropenem) for up to 14 days. Table 1 lists adverse reactions occurring in $\geq 1.0\%$ of patients receiving ZERBAXA[®] in Phase 3 clinical trials.

Table 1: Adverse Reactions Occurring in $\geq 1\%$ of Patients Receiving ZERBAXA[®] in Phase 3 Clinical Trials

Preferred Term	Complicated Intra-abdominal Infections		Complicated Urinary Tract Infections, Including Pyelonephritis	
	ZERBAXA [®] ^a (N=482) n (%)	Comparator ^b (N=497) n (%)	ZERBAXA [®] ^a (N=533) n (%)	Comparator ^c (N=535) n (%)
<i>Blood and the lymphatic system disorders</i>				
Anemia	7 (1.5)	5 (1)	2 (0.4)	5 (0.9)
Thrombocytosis	9 (1.9)	5 (1)	2 (0.4)	2 (0.4)
<i>Cardiac disorders</i>				
Atrial fibrillation	6 (1.2)	3 (0.6)	1 (0.2)	0
<i>Gastrointestinal disorders</i>				
Abdominal pain	6 (1.2)	2 (0.4)	4 (0.8)	2 (0.4)
Constipation	9 (1.9)	6 (1.2)	21 (3.9)	17 (3.2)
Diarrhea	30 (6.2)	25 (5)	10 (1.9)	23 (4.3)
Nausea	38 (7.9)	29 (5.8)	15 (2.8)	9 (1.7)
Vomiting	16 (3.3)	20 (4)	6 (1.1)	6 (1.1)
<i>General disorders and administration site conditions</i>				
Infusion Site Reactions	3 (0.6)	6 (1.2)	7 (1.3)	11 (2.1)
Pyrexia	27 (5.6)	20 (4)	9 (1.7)	5 (0.9)
<i>Investigations</i>				
ALT increased	7 (1.5)	5 (1)	9 (1.7)	5 (0.9)
AST increased	5 (1)	3 (0.6)	9 (1.7)	5 (0.9)
<i>Metabolism and nutrition disorders</i>				

Hypokalemia	16 (3.3)	10 (2)	4 (0.8)	2 (0.4)
<i>Nervous system disorders</i>				
Dizziness	4 (0.8)	5 (1)	6 (1.1)	1 (0.2)
Headache	12 (2.5)	9 (1.8)	31 (5.8)	26 (4.9)
<i>Psychiatric disorders</i>				
Anxiety	9 (1.9)	7 (1.4)	1 (0.2)	4 (0.7)
Insomnia	17 (3.5)	11 (2.2)	7 (1.3)	14 (2.6)
<i>Skin and subcutaneous tissue disorders</i>				
Rash	8 (1.7)	7 (1.4)	5 (0.9)	2 (0.4)
<i>Vascular disorders</i>				
Hypotension	8 (1.7)	4 (0.8)	2 (0.4)	1 (0.2)

^a The ZERBAXA[®] dose was 1.5 g IV every 8 hours, adjusted to match renal function where appropriate. In the complicated intra-abdominal infection studies ZERBAXA[®] was given in conjunction with metronidazole.

^b Meropenem 1 g every 8 hours

^c Levofloxacin 750 mg once daily

Less Common Clinical Trial Adverse Drug Reactions (< 1%)

The following selected adverse reactions were reported in ZERBAXA[®]-treated subjects at a rate of less than 1%:

Cardiac disorders: tachycardia, angina pectoris

Gastrointestinal disorders: gastritis, abdominal distension, dyspepsia, flatulence, ileus paralytic, *C. difficile* colitis

Infections and infestations: candidiasis including oropharyngeal and vulvovaginal, fungal urinary tract infection

Investigations: increased serum gamma-glutamyl transpeptidase (GGT), increased serum alkaline phosphatase, positive Coombs test

Metabolism and nutrition disorders: hyperglycemia, hypomagnesemia, hypophosphatemia

Nervous system disorders: ischemic stroke

Renal and urinary system: renal impairment, renal failure

Respiratory, thoracic and mediastinal disorders: dyspnea

Skin and subcutaneous tissue disorders: urticaria

Vascular disorders: venous thrombosis

Increased Mortality

In the cIAI trials (Phase 2 and 3), death occurred in 2.5% (14/564) of patients receiving ZERBAXA[®] and in 1.5% (8/536) of patients receiving meropenem. The causes of death varied and included worsening and/or complications of infection, surgery and underlying conditions.

DRUG INTERACTIONS

Overview

In vitro and *in vivo* data indicate that ZERBAXA[®] (ceftolozane and tazobactam) is unlikely to cause clinically relevant drug-drug interactions related to cytochrome P450 enzymes and transporters at therapeutic concentrations (see **ACTION AND CLINICAL PHARMACOLOGY**).

Drug-Drug Interactions

Tazobactam is a known substrate for OAT1 and OAT3 *in vitro*. Co-administration of ZERBAXA[®] with drugs that inhibit OAT1 and/or OAT3 (e.g., probenecid) may increase tazobactam plasma concentrations (see **ACTION AND CLINICAL PHARMACOLOGY**). No dose adjustment is recommended based on drug-drug interactions.

Drug-Food Interactions

Interactions with food have not been studied.

Drug-Herb Interactions

Interactions with herbal products have not been studied.

Drug-Laboratory Test Interactions

Interactions with laboratory tests have not been studied.

DOSAGE AND ADMINISTRATION**Dosing Considerations**

Because ZERBAXA[®] is eliminated primarily by the kidneys, a dosage adjustment is required for patients whose creatinine clearance is < 50 mL/min.

Recommended Dose and Dosage Adjustment

The recommended dosage regimen of ZERBAXA[®] is 1.5 g (1 g ceftolozane and 0.5 g tazobactam) administered every 8 hours by intravenous (IV) infusion over 1 hour in patients \geq 18 years of age. The duration of therapy should be guided by the severity and site of infection and the patient's clinical and bacteriological progress (Table 2).

Table 2: Dosage of ZERBAXA[®] by Infection in Patients with Creatinine Clearance (CrCL) >50 mL/min

Infection	Dose	Frequency	Infusion Time (hours)	Duration of Treatment
Complicated Intra-abdominal Infections*	1.5 g (1 g ceftolozane and 0.5 g tazobactam)	Every 8 Hours	1	4-14 days
Complicated Urinary Tract Infections, including Pyelonephritis	1.5 g (1 g ceftolozane and 0.5 g tazobactam)	Every 8 Hours	1	7 days

*In the treatment of cIAI, ZERBAXA[®] should be used in combination with metronidazole 500 mg intravenously every 8 hours in order to provide adequate anaerobic coverage.

Patients with Renal Impairment

ZERBAXA[®] is eliminated primarily by the kidneys; therefore a dosage adjustment is required for patients whose creatinine clearance is ≤ 50 mL/min, as shown in Table 3. For patients with changing renal function, monitor CrCl regularly and adjust the dosage of ZERBAXA[®] accordingly (see **WARNINGS AND PRECAUTIONS, ADVERSE REACTIONS, and ACTION AND CLINICAL PHARMACOLOGY**).

Table 3: Dosage of ZERBAXA[®] in Patients with Renal Impairment

Estimated CrCL (mL/min)*	Recommended Dosage Regimen for ZERBAXA [®] **
30 to 50	750 mg (500 mg ceftolozane and 250 mg tazobactam) intravenously every 8 hours
15 to 29	375 mg (250 mg ceftolozane and 125 mg tazobactam) intravenously every 8 hours
End stage renal disease (ESRD) on hemodialysis (HD)	A single loading dose of 750 mg (500 mg ceftolozane and 250 mg tazobactam) followed by a 150 mg (100 mg ceftolozane and 50 mg tazobactam) maintenance dose administered every 8 hours for the remainder of the treatment period (on hemodialysis days the dose should be administered at the earliest possible time following completion of dialysis)

*CrCL estimated using Cockcroft-Gault formula

**All doses of ZERBAXA[®] are administered over 1 hour

Missed Dose

If a dose is missed, it should be given as soon as possible. However, if it is less than two hours before the time for the next dose, no additional dose should be given and the regular dosing schedule should be resumed.

Preparation of Solutions

ZERBAXA[®] does not contain a bacteriostatic preservative. Aseptic technique must be followed in preparing the infusion solution.

Preparation of doses:

Reconstitute the vial with 10 mL of sterile water for injection or 0.9% Sodium Chloride for injection, USP (normal saline) and swirl to dissolve. The final volume is approximately 11.4 mL. The resultant concentration is approximately 132 mg/mL. **CAUTION: THE RECONSTITUTED SOLUTION IS NOT FOR DIRECT INJECTION.**

Table 4: Preparation of reconstituted solution

Diluent type	Diluent volume added to vial	Nominal total volume of the reconstituted solution	Nominal concentration of the reconstituted solution
Sterile Water for Injection	10 mL	11.4 mL	132 mg/mL (1.5 g/11.4mL)
0.9% Sodium Chloride for Injection, USP	10 mL	11.4 mL	
Note: Upon reconstitution, inspect the vial to ensure complete dissolution and ensure that the solution contains no particulate matter and no cake or powder remains attached to the sides of the vial.			

To prepare the required dose, using a syringe withdraw the appropriate volume (as determined from Table 5) from the reconstituted vial. Add the withdrawn volume to a Polyvinyl Chloride (PVC) infusion bag containing 100 mL of 0.9% Sodium Chloride for Injection, USP (normal saline) or 5% Dextrose Injection, USP. Use with Lactated Ringer's solution or 3.3% Dextrose and 0.3% Normal Saline has not been studied.

Table 5: Preparation of Doses

ZERBAXA [®] (ceftolozane and tazobactam) Dose	PVC IV bag containing infusion solution of	IV bag solution volume	Volume to withdraw from the reconstituted vial*	Total volume in IV bag (approximately)	Final concentration in IV bag (Nominal)
1.5 g (1 g ceftolozane and 0.5 g tazobactam)	0.9% Sodium chloride for Injection, USP	100 mL	~11.4 mL (entire contents)	~111.4 mL	13.5 mg/mL (1.5 g/111.4 mL)
750 mg (500 mg ceftolozane and 250 mg tazobactam)		100 mL	5.7 mL	105.7 mL	7 mg/mL (750 mg/105.7 mL)

375 mg (250 mg ceftolozane and 125 mg tazobactam)	Or	100 mL	2.9 mL	102.9 mL	4 mg/mL (375 mg/102.9 mL)
150 mg (100 mg ceftolozane and 50 mg tazobactam)	5% Dextrose	100 mL	1.2 mL	101.2 mL	1.6 mg/mL (150 mg/101.2 mL)

*Volumes for the withdrawal of the 750 mg, 375 and 150 mg doses are theoretical volumes based on the experimental extractable volume of entire vial (11.4 mL).

Prior to administration, parenteral drug products should be inspected for clarity, particulate matter, precipitate and discoloration. Solutions showing haziness, particulate matter, precipitate, discoloration should not be used. ZERBAXA[®] infusions range from clear, colorless solutions to solutions that are clear and slightly yellow. Variations in color within this range do not affect the potency of the product.

Compatibility

Compatibility of ZERBAXA[®] with other drugs has not been established. ZERBAXA[®] should not be mixed with other drugs or physically added to intravenous solutions containing other drugs.

Storage of Constituted Solutions

Upon reconstitution with sterile water for injection or 0.9% sodium chloride (normal saline) injection, ZERBAXA[®] solution in the vial may be held for 1 hour prior to transfer and dilution in the infusion bag at room temperature [(15-30°C)].

Following dilution of the solution with normal saline or 5% dextrose, ZERBAXA[®] is stable for 24 hours when stored at room temperature [(15-30°C)] or 7 days when stored under refrigeration at 2 to 8°C. Discard unused portions.

Reconstituted ZERBAXA[®] solution or ZERBAXA[®] infusion should not be frozen.

OVERDOSAGE

In the event of overdose, discontinue ZERBAXA[®] and provide general supportive treatment. ZERBAXA[®] can be removed by hemodialysis. Approximately 66% of ceftolozane, 56% of tazobactam, and 51% of the tazobactam metabolite M1 are removed by dialysis. However, no information is available on the use of hemodialysis to treat overdose. The highest single dose of ZERBAXA[®] received in clinical trials was 4.5 g (comprised of 3 g of ceftolozane and 1.5 g of tazobactam); at this dosage no adverse pharmacological effects or increased safety risks have been observed.

For management of a suspected drug overdose, contact your regional Poison Control Centre.

ACTION AND CLINICAL PHARMACOLOGY

Mechanism of Action

Ceftolozane is a cephalosporin class antibacterial drug. The bactericidal action of ceftolozane results from inhibition of cell wall biosynthesis, and is mediated through binding to penicillin-binding proteins (PBPs). Ceftolozane is an inhibitor of PBPs of *P. aeruginosa* (e.g., PBP1b, PBP1c, and PBP3) and *E. coli* (e.g., PBP3).

Tazobactam sodium has little clinically relevant *in vitro* activity against bacteria due to its reduced affinity to penicillin-binding proteins. It is an irreversible inhibitor of some beta lactamases (e.g., certain penicillinases and cephalosporinases), and can bind covalently to some chromosomal and plasmid-mediated bacterial beta-lactamases (see **MICROBIOLOGY**).

Pharmacodynamics

For ceftolozane, the time that the plasma concentration of ceftolozane exceeds the minimum inhibitory concentration (MIC) of the infecting organism has been shown to be the best predictor of efficacy in animal models of infection. For tazobactam, the time above a threshold concentration for the infecting organism has been determined to be the parameter that best predicts the efficacy of tazobactam in *in vitro* nonclinical models. The exposure-response analyses in Phase 2 trials support the recommended dose of ZERBAXA[®].

Cardiac Electrophysiology

In a randomized, positive and placebo-controlled crossover thorough QTc study, 51 healthy subjects were administered a single therapeutic dose (1.5 g) and a suprathreshold dose (4.5 g) of ceftolozane and tazobactam. No significant effects of ceftolozane and tazobactam on heart rate, electrocardiogram morphology, PR, QRS, or QT interval were detected. Therefore, ceftolozane and tazobactam does not affect cardiac repolarization.

Pharmacokinetics

The mean pharmacokinetic parameters of ceftolozane and tazobactam in healthy adults with normal renal function after single and multiple 1-hour IV infusions of 1.5 g ceftolozane and tazobactam administered every 8 hours are summarized in Table 6. Pharmacokinetic parameters were similar for single and multiple dose administration.

Table 6: Mean (CV%) Plasma Pharmacokinetic Parameters of ZERBAXA[®] (ceftolozane and tazobactam) After Single and Multiple Intravenous 1-hour Infusions of ZERBAXA[®] 1.5 g (ceftolozane 1 g and tazobactam 0.5 g) Every 8 Hours in Healthy Adults

PK parameters	ZERBAXA [®] 1.5 g (ceftolozane 1 g and tazobactam 0.5 g) (every 8 hours)			
	Ceftolozane		Tazobactam	
	Day 1 (n=9) ^a	Day 10 (n=10)	Day 1 (n=9) ^a	Day 10 (n=10)
C _{max} (µg/mL)	69.1 (11)	74.4 (14)	18.4 (16)	18.0 (8)
t _{max} (h) ^b	1.02 (1.01, 1.1)	1.07 (1.0, 1.1)	1.02 (0.99, 1.03)	1.01 (1.0, 1.1)
AUC (µg•h/mL) ^c	172 (14)	182 (15)	24.4 (18)	25.0 (15)
t _{1/2} (h)	2.77 (30)	3.12 (22)	0.91 (26) ^d	1.03 (19)

^a N = 9, one outlier subject excluded from descriptive statistics

^b Median (minimum, maximum) presented

^c AUC for Day 1 = AUC_{last} and AUC for Day 10 = steady state AUC (AUC_{τ,ss})

^d N = 8, one subject excluded from descriptive statistics as the concentration-time profile did not exhibit a terminal log-linear phase and t_{1/2} could not be calculated

The C_{max} and AUC of ceftolozane and tazobactam increase in proportion to dose. Plasma levels of ZERBAXA[®] do not increase appreciably following multiple IV infusions of up to 3.0 g (ceftolozane 2 g and tazobactam 1 g) administered every 8 hours for up to 10 days in healthy adults with normal renal function. The elimination half-life (t_{1/2}) of ceftolozane is independent of dose.

Distribution: The binding of ceftolozane and tazobactam to human plasma proteins is approximately 16% to 21% and 30%, respectively. The mean (CV%) steady-state volume of distribution of ZERBAXA[®] in healthy adult males (n = 51) following a single intravenous dose of ZERBAXA[®] 1.5 g (ceftolozane 1 g and tazobactam 0.5 g) was 13.5 L (21%) and 18.2 L (25%) for ceftolozane and tazobactam, respectively, similar to extracellular fluid volume. In rat and dog repeat-dose studies, ceftolozane accumulated in renal tissue with an estimated renal tissue elimination half-life of 76 hour in rats.

Metabolism: Ceftolozane is eliminated in the urine as unchanged parent drug and thus does not appear to be metabolized to any appreciable extent. The beta-lactam ring of tazobactam is hydrolyzed to form the pharmacologically inactive tazobactam metabolite, M1.

Excretion: Ceftolozane, tazobactam and the tazobactam metabolite M1 are eliminated by the kidneys. Following administration of a single ZERBAXA[®] 1.5 g (ceftolozane 1 g and tazobactam 0.5 g) intravenous dose to healthy male adults greater than 95% of ceftolozane was excreted in the urine as unchanged parent drug. More than 80% of tazobactam was excreted as the parent compound with the remainder excreted as the tazobactam M1 metabolite. After a single dose of ZERBAXA[®], renal clearance (CL) of ceftolozane (3.41 – 6.69 L/h) was similar to

plasma CL (4.10 to 6.73 L/h) and similar to the glomerular filtration rate for the unbound fraction, suggesting that ceftolozane is eliminated by the kidney via glomerular filtration.

Special Populations and Conditions

Pediatrics: Safety and effectiveness in pediatric patients have not been established.

Geriatrics: In a population pharmacokinetic analysis of ceftolozane and tazobactam, no clinically relevant trend in exposure was observed with regard to age.

Dosage adjustment for elderly patients should be based on renal function.

Gender: In a population pharmacokinetic analysis of ceftolozane and tazobactam, no clinically relevant differences in AUC were observed for ceftolozane (116 males compared to 70 females) or tazobactam (80 males compared to 50 females).

No dose adjustment is recommended based on gender.

Race: In a population pharmacokinetic analysis of ceftolozane and tazobactam, no clinically relevant differences in ceftolozane and tazobactam AUC were observed in Caucasians (n = 156) compared to all other races combined (n = 30).

No dose adjustment is recommended based on race.

Hepatic Insufficiency: As ceftolozane and tazobactam does not undergo hepatic metabolism, the systemic clearance of ZERBAXA[®] is not expected to be affected by hepatic impairment.

No dose adjustment is recommended for ZERBAXA[®] in subjects with hepatic impairment.

Renal Insufficiency: Ceftolozane, tazobactam and the tazobactam metabolite M1 are eliminated by the kidneys.

The ceftolozane dose normalized geometric mean AUC increased up to 1.26-fold, 2.5-fold, and 5-fold in subjects with mild, moderate, and severe renal impairment, respectively, compared to healthy subjects with normal renal function. The respective tazobactam dose normalized geometric mean AUC increased approximately up to 1.3-fold, 2-fold, and 4-fold. To maintain similar systemic exposures to those with normal renal function, dosage adjustment is required (see **DOSAGE AND ADMINISTRATION**).

In subjects with end-stage renal disease (ESRD) on hemodialysis (HD), approximately two-thirds of the administered ceftolozane and tazobactam dose is removed by HD. The recommended dose in subjects with ESRD on HD is a single loading dose of 750 mg ceftolozane and tazobactam followed by a 150 mg maintenance dose of ceftolozane and tazobactam administered every 8 hours for the remainder of the treatment period. On HD days, the dose should be administered at the earliest possible time following completion of HD (see **DOSAGE AND ADMINISTRATION**).

Drug Interactions

No drug-drug interaction was observed between ceftolozane and tazobactam in a clinical study in 15 healthy subjects. *In vitro* and *in vivo* data indicate that ZERBAXA[®] is unlikely to cause clinically relevant drug-drug interactions related to CYPs and transporters at therapeutic concentrations.

Drug Metabolizing Enzymes

In vivo data indicated that ZERBAXA[®] is not a substrate for CYPs. Thus clinically relevant drug-drug interactions involving inhibition or induction of CYPs by other drugs are unlikely to occur.

In vitro studies demonstrated that ceftolozane, tazobactam and the M1 metabolite of tazobactam did not inhibit CYP1A2, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP2D6, or CYP3A4 and did not induce CYP1A2, CYP2B6, or CYP3A4 at therapeutic plasma concentrations. *In vitro* induction studies in primary human hepatocytes demonstrated that ceftolozane, tazobactam, and the tazobactam metabolite M1 decreased CYP1A2 and CYP2B6 enzyme activity and mRNA levels in primary human hepatocytes as well as CYP3A4 mRNA levels at supratherapeutic plasma concentrations (at concentrations that were 17.5-fold, 56.8-fold, and 20-fold higher than the mean clinical C_{max} for ceftolozane (57 mcg/mL), tazobactam (22 mcg/mL), and tazobactam metabolite M1 (1.5 mcg/mL), respectively, in cIAI patients with normal renal function following administration of 1 g ceftolozane and 0.5 g tazobactam). Tazobactam metabolite M1 also decreased CYP3A4 activity at supratherapeutic plasma concentrations. A clinical drug-drug interaction study was conducted and results indicated drug interactions involving CYP1A2 and CYP3A4 inhibition by ZERBAXA[®] are not anticipated.

Membrane Transporters

Ceftolozane and tazobactam were not substrates for P-gp or BCRP, and tazobactam was not a substrate for OCT2, *in vitro* at therapeutic concentrations.

Tazobactam is a known substrate for OAT1 and OAT3. Co-administration of tazobactam with OAT1/OAT3 inhibitor probenecid has been shown to prolong the half-life of tazobactam by 71%. Co-administration of ZERBAXA[®] with drugs that inhibit OAT1 and/or OAT3 may increase tazobactam plasma concentrations.

In vitro data indicate that ceftolozane did not inhibit P-gp, BCRP, OATP1B1, OATP1B3, OCT1, OCT2, MRP2, BSEP, OAT1, OAT3, MATE1, or MATE2-K *in vitro* at therapeutic plasma concentrations.

In vitro data indicate that neither tazobactam nor the tazobactam metabolite M1 inhibit P-gp, BCRP, OATP1B1, OATP1B3, OCT1, OCT2, or BSEP transporters at therapeutic plasma concentrations. *In vitro*, tazobactam, but not tazobactam M1 metabolite, inhibited human OAT1 and OAT3 transporters with IC₅₀ values of 118 and 147 ug/mL, respectively. A clinical drug-drug interaction study was conducted and results indicated drug interactions involving OAT1/OAT3 inhibition by ZERBAXA[®] are not anticipated.

Enzyme Induction:

Hepatic mixed function oxidase studies in the rat and dog indicated that tazobactam did not induce the hepatic drug metabolizing enzymes in these species. No enzyme induction studies were conducted in rats or dogs with ceftolozane.

STORAGE AND STABILITY

ZERBAXA[®] vials should be stored under refrigeration at 2 to 8°C. Protect from exposure to light.

DOSAGE FORMS, COMPOSITION AND PACKAGING

ZERBAXA[®] (ceftolozane and tazobactam) for injection is a white to yellow sterile lyophilized powder for reconstitution, supplied in single-dose, Type 1, 20 mL clear glass vial with a bromobutyl, siliconized stopper and a purple flip-off seal. Each vial contains 1 gram ceftolozane (as ceftolozane sulfate) and 500 mg tazobactam (as tazobactam sodium), and 10 vials are packaged in each carton.

The product contains sodium chloride, citric acid, and L-arginine.

PART II: SCIENTIFIC INFORMATION

PHARMACEUTICAL INFORMATION


Drug Substance

Proper name: Ceftolozane Sulfate

Chemical name: 1*H*-Pyrazolium, 5-amino-4-[[[(2-aminoethyl)amino]carbonyl]amino]-2-[[[(6*R*,7*R*)-7-[[[(2*Z*)-2-(5-amino-1,2,4-thiadiazol-3-yl)-2-[(1-carboxy-1-methylethoxy)imino]acetyl]amino]-2-carboxy-8-oxo-5-thia-1-azabicyclo[4.2.0]oct-2-en-3-yl]methyl]-1-methyl-,sulfate (1:1).

Molecular formula and molecular mass: $C_{23}H_{31}N_{12}O_8S_2^+ \cdot HSO_4^-$ MW = 764.77 g/mol

Structural formula:



Physicochemical properties: Ceftolozane sulfate is a white to off-white powder that is hygroscopic. It is slightly soluble in water and insoluble in isopropanol. The pH in a 2% aqueous solution is 1.92. The optical rotation, $[\alpha]_D^{20}$ of a 10.144 mg/mL solution of ceftolozane sulfate in 1N aqueous HCl was -14.9° .


Drug Substance

Proper name: Tazobactam Sodium

Chemical name: sodium (2*S*,3*S*,5*R*)-3-methyl-7-oxo-3-(1*H*-1,2,3-triazol-1-ylmethyl)-4-thia-1-azabicyclo[3.2.0]heptane-2-carboxylate-4,4-dioxide

Molecular formula and molecular mass: $C_{10}H_{11}N_4NaO_5S$ MW = 322.3 g/mole

Structural formula:



Physicochemical properties: Sterile tazobactam sodium drug substance is a white to off-white powder that is hygroscopic. It is freely soluble in water and slightly soluble in ethanol and acetone. The pH of an aqueous solution of the drug substance is 5.7 – 6.7. The specific optical rotation of the drug substance ranges between +138.0° and +152.0°.

ZERBAXA[®], ceftolozane and tazobactam parenteral combination, is a white to yellow sterile powder consisting of ceftolozane sulfate (1147 mg/vial equivalent to 1 g of ceftolozane) and tazobactam sodium (537 mg/vial equivalent to 0.5 g of tazobactam).

CLINICAL TRIALS

Study demographics and trial design

The trial design and patient demographics for the ZERBAXA[®] clinical trials are summarized in Table 7 below.

Table 7 Summary of patient demographics for Phase 3 clinical trials in specific indications^{1,2}

Study #	Trial design	Dosage, route of administration and duration	Number of Subjects	Mean age (Range)	Gender
Trial 1 CXA-cIAI-10-08 and CXA-cIAI-10-09 (complicated intra-abdominal infections)	Randomized, placebo-controlled, double-blind, multicentered	ZERBAXA [®] 1.5 g every 8 hours and metronidazole 500 mg every 8 hours, 1-hour IV infusion Meropenem 1 g every 8 hours, 1-hour IV infusion 4 to 10 days	806	51 years (18 to 92 years)	Male: 56% Female: 44%

Study #	Trial design	Dosage, route of administration and duration	Number of Subjects	Mean age (Range)	Gender
Trial 2 CXA-cUTI-10-04 and CXA-cUTI-10-05 (complicated urinary tract infection, including pyelonephritis)	Randomized, double-blind, multicentered	ZERBAXA [®] (1.5 g every 8 hours), 1-hour IV infusion versus levofloxacin (750 mg once daily), 1-hour IV infusion 7 days	800	48 years (31 to 65 years)	Male: 26% Female: 74%

Study results

Complicated Intra-abdominal Infections

The primary efficacy endpoint was clinical response at the test-of-cure (TOC) visit in the microbiological intent-to-treat (MITT) population, which included all patients who had at least 1 baseline intra-abdominal pathogen. The key secondary efficacy endpoint was clinical response at the TOC visit in the microbiologically evaluable (ME) population, which included all protocol-adherent MITT patients.

Clinical cure rates at the TOC visit are displayed by patient population in Table 8. Clinical cure rates at the TOC visit by pathogen in the ME population are presented in Table 9.

Table 8: Clinical Cure Rates in a Phase 3 Study of Complicated Intra-Abdominal Infections

Analysis Population	ZERBAXA [®] plus metronidazole ^a n/N (%)	Comparator ^b n/N (%)	Treatment Difference (95% CI) ^c
MITT	323/389 (83.0)	364/417 (87.3)	-4.2 (-8.91, 0.54)
ME	259/275 (94.2)	304/321 (94.7)	-1.0 (-4.52, 2.59)

^a ZERBAXA[®] 1.5 g IV every 8 hours + metronidazole 500 mg IV every 8 hours

^b Meropenem 1 g IV every 8 hours

^c The 95% CI was calculated using the Newcombe method with minimum risk weights

Table 9: Clinical Cure Rates by Pathogen in a Phase 3 Study of Complicated Intra-abdominal Infections (ME Population)

Organism Group Pathogen	ZERBAXA [®] plus metronidazole n/N (%)	Comparator ^a n/N (%)
Aerobic Gram-negative	238/252 (94.4)	273/291 (93.8)
<i>Escherichia coli</i>	197/208 (94.7)	216/231 (93.5)

Organism Group Pathogen	ZERBAXA [®] plus metronidazole n/N (%)	Comparator ^a n/N (%)
<i>Klebsiella pneumoniae</i>	28/30 (93.3)	22/25 (88.0)
<i>Pseudomonas aeruginosa</i>	26/26 (100)	27/29 (93.1)
<i>Enterobacter cloacae</i>	19/22 (86.4)	22/22 (100)
<i>Klebsiella oxytoca</i>	12/12 (100)	21/22 (95.5)
<i>Proteus mirabilis</i>	10/11 (90.9)	9/10 (90.0)
Aerobic Gram-positive	153/168 (91.1)	170/185 (91.9)
<i>Streptococcus anginosus</i>	25/30 (83.3)	23/23 (100)
<i>Streptococcus constellatus</i>	17/18 (94.4)	20/23 (87.0)
<i>Streptococcus salivarius</i>	9/10 (90.0)	8/8 (100)
Anaerobic Gram-negative	104/109 (95.4)	132/137 (96.4)
<i>Bacteroides fragilis</i>	39/41 (95.1)	56/57 (98.2)
<i>Bacteroides ovatus</i>	36/37 (97.3)	42/42 (100)
<i>Bacteroides thetaiotaomicron</i>	20/20 (100)	40/43 (93.0)
<i>Bacteroides vulgatus</i>	12/13 (92.3)	21/22 (95.5)

^a Meropenem 1 g IV every 8 hours

In a subset of the *E. coli* and *K. pneumoniae* isolates from both arms of the cIAI Phase 3 trial that met pre-specified criteria for beta-lactam susceptibility, genotypic testing identified certain ESBL groups (e.g., TEM, SHV, CTX-M, OXA) in 53/601 (9%). Cure rates in this subset were similar to the overall trial results. *In vitro* susceptibility testing showed that some of these isolates were susceptible to ZERBAXA[®] (MIC ≤ 2 mcg/mL), while some others were not susceptible (MIC > 2 mcg/mL). Isolates of a specific genotype were seen in patients who were deemed to be either successes or failures.

Complicated Urinary Tract Infections, including Pyelonephritis

The primary efficacy endpoint was defined as complete resolution or marked improvement of the clinical symptoms and microbiological eradication (all uropathogens found at baseline at $\geq 10^5$ were reduced to $< 10^4$ CFU/mL) at the test-of-cure (TOC) visit 7 (± 2) days after the last dose of study drug. The primary efficacy analysis population was the microbiologically modified intent-to-treat (mMITT) population, which included all patients who received study medication and had at least 1 baseline uropathogen. The key secondary efficacy endpoint was the composite microbiological and clinical cure response at the TOC visit in the microbiologically evaluable (ME) population, which included protocol-adherent mMITT patients with a urine culture at the TOC visit.

Concomitant bacteremia was identified in 62 (7.8%) patients at baseline (ITT population). In the mMITT population, composite cure rates in patients with concurrent bacteremia were 23/29 (79.3%) for ZERBAXA[®].

Composite microbiological and clinical cure rates at the TOC visit in both the mMITT and ME populations are shown in Table 10. Microbiological eradication rates at the TOC visit by pathogen in the ME population are presented in Table 11.

Although a statistically significant difference was observed in the ZERBAXA[®] arm compared to the levofloxacin arm with respect to the primary endpoint, it was likely attributable to the 212/800 (26.5%) patients with baseline organisms non-susceptible to levofloxacin. Among

patients infected with a levofloxacin-susceptible organism at baseline, the response rates were similar between treatment arms (Table 10).

Table 10: Composite Microbiological and Clinical Cure Rates in a Phase 3 Trial of Complicated Urinary Tract Infections

Analysis Population	ZERBAXA [®] ^a n/N (%)	Comparator ^b n/N (%)	Treatment Difference (95% CI) ^c
mMITT	306/398 (76.9)	275/402 (68.4)	8.5 (2.3, 14.6)
Levofloxacin resistant baseline pathogen(s)	60/100 (60)	44/112 (39.3)	
No levofloxacin resistant baseline pathogen(s)	246/298 (82.6)	231/290 (79.7)	
ME	284/341 (83.3)	266/353 (75.4)	8.0 (2.0, 14.0)

^a ZERBAXA[®] 1.5 g intravenously every 8 hours

^b Levofloxacin 750 mg intravenously once daily

^c The 95% confidence interval was based on the stratified Newcombe method.

Table 11: Per Pathogen Microbiological Eradication Rates in a Phase 3 Study of Complicated Urinary Tract Infections (ME Population)

Organism Group Pathogen	ZERBAXA [®] n/N (%)	Comparator ^a n/N (%)
Aerobic Gram-negative	287/323 (88.9)	263/340 (77.4)
<i>Escherichia coli</i>	237/262 (90.5)	226/284 (79.6)
<i>Klebsiella pneumoniae</i>	21/25 (84.0)	14/23 (60.9)
<i>Proteus mirabilis</i>	10/10 (100)	8/11 (72.7)
<i>Pseudomonas aeruginosa</i>	6/7 (85.7)	7/12 (58.3)

^a Levofloxacin 750 mg intravenously once daily

In a subset of the *E. coli* and *K. pneumoniae* isolates from both arms of the cUTI Phase 3 trial that met pre-specified criteria for beta-lactam susceptibility, genotypic testing identified certain ESBL groups (e.g., TEM, SHV, CTX-M, OXA) in 104/687 (15%). Cure rates in this subset were similar to the overall trial results. *In vitro* susceptibility testing showed that some of these isolates were susceptible to ZERBAXA[®] (MIC ≤ 2 mcg/mL), while some others were not susceptible (MIC > 2 mcg/mL). Isolates of a specific genotype were seen in patients who were deemed to be either successes or failures.

MICROBIOLOGY

Mechanism of Action

See ACTION AND CLINICAL PHARMACOLOGY.

Mechanism(s) of Resistance

Mechanisms of beta-lactam resistance may include the production of beta-lactamases, modification of PBPs by gene acquisition or target alteration, up-regulation of efflux pumps, and loss of outer membrane porin.

Clinical isolates may produce multiple beta-lactamases, express varying levels of beta-lactamases, or have amino acid sequence variations, and other resistance mechanisms that have not been identified.

Culture and susceptibility information and local epidemiology should be considered in selecting or modifying antibacterial therapy.

ZERBAXA[®] demonstrated *in vitro* activity against Enterobacteriaceae in the presence of some extended-spectrum beta-lactamases (ESBLs) and other beta-lactamases of the following groups: TEM, SHV, CTX-M, and OXA. ZERBAXA[®] is not active against bacteria that produce serine carbapenamases [*K. pneumoniae* carbapenemase (KPC)], and metallo-beta lactamases.

In ZERBAXA[®] clinical trials, some isolates of *E. coli* and *K. pneumoniae*, that produced beta-lactamases, were susceptible to ZERBAXA[®] (minimum inhibitory concentration ≤ 2 mcg/mL). These isolates produced one or more beta-lactamases of the following enzyme groups: CTX-M, OXA, TEM, or SHV.

Some of these beta-lactamases were also produced by isolates of *E. coli* and *K. pneumoniae* that were not susceptible to ZERBAXA[®] (minimum inhibitory concentration > 2 mcg/mL). These isolates produced one or more beta-lactamases of the following enzyme groups: CTX-M, OXA, TEM, or SHV.

ZERBAXA[®] demonstrated *in vitro* activity against *P. aeruginosa* isolates tested that had chromosomal AmpC, loss of outer membrane porin (OprD), or up-regulation of efflux pumps (MexXY, MexAB).

Cross-Resistance

Isolates resistant to other cephalosporins may be susceptible to ceftolozane and tazobactam, although cross-resistance may occur.

Interaction with Other Antimicrobials

In vitro synergy studies suggest that ceftolozane and tazobactam has negligible potential to antagonize or be antagonized by other antibiotics (e.g. meropenem, amikacin, aztreonam, levofloxacin, tigecycline rifampin, linezolid, daptomycin, vancomycin, and metronidazole).

List of Microorganisms

ZERBAXA[®] has been shown to be active against the following bacteria, both *in vitro* and in clinical infections (see **INDICATIONS AND CLINICAL USE**).

Complicated Intra-abdominal Infections

Gram-negative bacteria

Enterobacter cloacae

Escherichia coli

Klebsiella oxytoca

Klebsiella pneumoniae
Proteus mirabilis
Pseudomonas aeruginosa

Gram-positive bacteria

Streptococcus anginosus
Streptococcus constellatus
Streptococcus salivarius

Gram-negative Anaerobes

Bacteroides fragilis

Complicated Urinary Tract Infections, Including Pyelonephritis

Gram-negative bacteria

Escherichia coli
Klebsiella pneumoniae
Proteus mirabilis
Pseudomonas aeruginosa

The following *in vitro* data are available, **but their clinical significance is unknown.** At least 90% of isolates of the following bacteria, in the absence of acquired mechanisms of resistance, exhibited an *in vitro* minimum inhibitory concentration (MIC) less than or equal to 2/4 mcg/mL ceftolozane/tazobactam. The safety and effectiveness of ZERBAXA[®] in treating clinical infections due to these bacteria have not been established.

Gram-negative bacteria

Citrobacter koseri
Haemophilus influenzae
Moraxella catarrhalis
Morganella morganii
Proteus vulgaris
Providencia rettgeri
Providencia stuartii
Serratia liquefaciens
Serratia marcescens

Gram-positive aerobic bacteria

Streptococcus agalactiae
Streptococcus intermedius
Streptococcus pyogenes
Streptococcus pneumoniae

In vitro data indicate that the following bacteria are not susceptible to ceftolozane and tazobactam:

Staphylococcus aureus
Enterococcus faecalis
Enterococcus faecium

Susceptibility Test Methods

When available, the clinical microbiology laboratory should provide the results of *in vitro* susceptibility testing for antimicrobial drugs used in local hospitals and practice areas to the physician as periodic reports that describe the susceptibility profile of nosocomial and community-acquired pathogens. These reports should aid the physician in selecting an antibacterial drug product for treatment.

Dilution Techniques:

Quantitative methods are used to determine antibacterial MICs. Ceftolozane and tazobactam susceptibility testing is performed with a fixed 4 mcg/mL concentration of tazobactam. These MICs provide estimates of the susceptibility of bacteria to antibacterial compounds. The MICs should be determined using a standardized test method (broth, and/or agar).^{3,6} The MIC values should be interpreted according to the criteria in Table 12.

Diffusion Techniques:

Quantitative methods that require measurement of zone diameters can also provide reproducible estimates of the susceptibility of bacteria to antibacterial compounds. The zone size should be determined using a standardized test method.^{4,6} This procedure uses paper disks impregnated with ceftolozane 30 mcg and tazobactam 10 mcg to test the susceptibility of bacteria to ceftolozane and tazobactam. Results should be interpreted according to the criteria in Table 12.

Anaerobic Techniques:

For anaerobic bacteria, the susceptibility to ceftolozane and tazobactam should be determined using a standardized test method.⁵ The MIC values obtained should be interpreted according to criteria provided in Table 12.

Table 12: Susceptibility Interpretive Criteria for Ceftolozane and Tazobactam

Pathogen and Isolate Source	Minimum Inhibitory Concentrations (mcg/mL) (ceftolozane/tazobactam)			Disk Diffusion Zone Diameter (mm)		
	S	I	R	S	I	R
Enterobacteriaceae	≤2/4	4/4	≥8/4	≥21	18-20	≤17
<i>Pseudomonas aeruginosa</i>	≤4/4	8/4	≥16/4	≥21	17-20	≤16
<i>Streptococcus anginosus</i> <i>Streptococcus constellatus</i> and <i>Streptococcus salivarius</i>	≤8/4	16/4	≥32/4	---	---	---
<i>Bacteroides fragilis</i>	≤8/4	16/4	≥32/4	---	---	---

S = susceptible, I = intermediate, R = resistant

A report of “Susceptible” indicates that the antibacterial is likely to inhibit growth of the bacteria if the antibacterial drug reaches the concentration usually achievable at the site of infection. A report of “Intermediate” indicates that the result should be considered equivocal, and if the bacteria is not fully susceptible to alternative clinically feasible drugs, the test should be repeated. This category implies possible clinical applicability in body sites where the drug is physiologically concentrated. This category also provides a buffer zone that prevents small uncontrolled technical factors from causing major discrepancies in interpretation. A report of “Resistant” indicates that the antibacterial is not likely to inhibit growth of the bacteria if the antibacterial drug reaches the concentrations usually achievable at the infection site; other therapy should be selected.

Quality Control

Standardized susceptibility test procedures require the use of laboratory controls to monitor and ensure the accuracy and precision of supplies and reagents used in the assay, and the techniques of the individuals performing the test.^{3,4,5,6} Standard ceftolozane and tazobactam powder should provide the following range of MIC values provided in Table 13. For the diffusion technique using the ceftolozane 30 mcg and tazobactam 10 mcg disk, the criteria provided in Table 13 should be achieved.⁶

Table 13: Acceptable Quality Control Ranges for Ceftolozane and Tazobactam

Quality Control Organism	Minimum Inhibitory Concentrations (mcg/mL) (ceftolozane/tazobactam)	Disk Diffusion Zone Diameters (mm)
<i>Escherichia coli</i> ATCC 25922	0.12/4-0.5/4	24-32
<i>Escherichia coli</i> ^a ATCC 35218	0.06/4-0.25/4	25-31
<i>Pseudomonas aeruginosa</i> ATCC 27853	0.25/4-1/4	25-31
<i>Staphylococcus aureus</i> ATCC 25923	Not applicable	10-18
<i>Staphylococcus aureus</i> ATCC 29213	16/4-64/4	Not Applicable
<i>Haemophilus influenzae</i> ^b ATCC 49247	0.5/4-2/4	23-29
<i>Klebsiella pneumoniae</i> ^a ATCC 700603	0.5/4-2/4	17-25
<i>Streptococcus pneumoniae</i> ATCC 49619	0.25/4-1/4	21-29
<i>Bacteroides fragilis</i> ATCC 25285 (agar and broth)	0.12/4-1/4	Not Applicable
<i>Bacteroides thetaiotaomicron</i> ATCC 29741 (agar)	16/4-128/4	Not Applicable
<i>Bacteroides thetaiotaomicron</i> ATCC 29741 (broth)	16/4-64/4	Not Applicable

ATCC = American Type Culture Collection

^a Store *E. coli* ATCC 35218 and *K. pneumoniae* ATCC 700603 stock cultures at -60°C or below and prepare working stock cultures weekly.

^bThis strain may lose its plasmid and develop susceptibility to beta-lactam antimicrobial agents after repeated transfers onto culture media. Minimize by removing new culture from storage at least monthly or whenever the strain begins to show increased zone diameters to ampicillin, piperacillin, or ticarcillin.

TOXICOLOGY

Repeat-Dose Toxicity

Ceftolozane

No clinically-relevant adverse effects were evident in adult rats and dogs up to the highest ceftolozane IV doses (1000 mg/kg/day) tested in repeat dose toxicity studies (up to 28 days). Microscopic evidence of hyaline droplets (confirmed as secondary lysosomes by electron microscopy) were detected in proximal renal tubules of rats and dogs following once daily repeated IV administration of ceftolozane for 28 days at doses of 300 and 1000 mg/kg/day. A corresponding increase in kidney weight was consistently observed at these dose levels in rats but not dogs. No microscopic evidence of renal tubular degeneration or necrosis was detected and no toxicologically relevant effect on renal function was noted as determined by biologically relevant changes in serum blood urea nitrogen (BUN), creatinine, inorganic phosphorus, or urine volume. In both rats and dogs, the ceftolozane-related effects on the kidney had reversed or were in the process of reversing following a 28 day recovery period. NOAELs in adult rats and dogs were considered 1000 and 300 mg/kg/day, respectively. The NOAEL for dogs (300 mg/kg/day) was based on the presence of adverse ceftolozane-induced histamine-related clinical signs at 1000 mg/kg/day; namely flush of auricles and oral mucosa, swelling of the head, emesis, salivation, and lateral (prone) position. Ceftolozane AUC and C_{max} values at the NOAEL are, respectively, at least ~3-fold and 13-fold greater than the clinical AUC and C_{max} values at the therapeutic dose (1000 mg thrice daily).

Tazobactam

Repeat-dose toxicity studies of up to 6 months in the rat and dog showed reversible hepatocellular glycogen accumulation in rats at intraperitoneal doses ≥ 80 mg/kg/day and in dogs at intravenous doses ≥ 80 mg/kg/day. Other drug-related effects observed in rats were decreased red blood cell parameters and decreased serum triglycerides and enlarged ceca. The effects on red blood cell parameters and triglyceride levels were reversible. The NOAELs determined for both species were 40 mg/kg/day.

Ceftolozane and Tazobactam

No new effects or unexpected or altered toxicities were identified when ceftolozane and tazobactam were administered intravenously to adult rats for 28 days or to dogs for 14 days, in a fixed 2:1 ratio as compared to the individual agents. The NOAEL for ceftolozane and tazobactam was considered 1000 and 500 mg/kg/day in rats and 300 and 150 mg/kg/day in dogs, respectively. Ceftolozane AUC and C_{max} values at the NOAELs are, respectively, at least approximately 3-fold and 16-fold greater than the clinical AUC and C_{max} values at the therapeutic dose (1g thrice daily). Tazobactam AUC and C_{max} values at the NOAELs are, respectively, at least approximately 1-fold and 10-fold greater than the clinical AUC and C_{max} values at the therapeutic dose (500 mg thrice daily).

In neonatal rats dosed subcutaneously with ceftolozane and tazobactam (1000 and 500 mg/kg/day) from Postnatal Day 4 (PND4) to 17, minimal to mild renal tubular basophilia and cortical fibrosis were observed (plasma exposures approximately 13-fold and 5-fold the clinical AUC on PND4 and 17, respectively). At the NOAEL (300 and 150 mg/kg/day), the safety margins based on AUC were 3.2 and 1.4-fold, respectively. Similar effects were not observed in a longer study in neonatal rats (PND4-31) at the same doses.

Genotoxicity

The genotoxic potential of ceftolozane, alone and in combination with tazobactam, was characterized in *in silico*, *in vitro* and *in vivo* genotoxicity studies. ZERBAXA[®] was not genotoxic *in vivo*.

Ceftolozane

Ceftolozane was negative for mutagenic potential in an *in silico* screen. Ceftolozane was negative for genotoxicity in an *in vitro* microbial mutagenicity (Ames) assay, an *in vitro* chromosomal aberration assay in Chinese hamster lung fibroblast cells, an *in vitro* mouse lymphoma assay, an *in vitro* hypoxanthine-guanine phosphoribosyl transferase (HPRT) assay in Chinese hamster ovary cells, an *in vivo* mouse micronucleus assay, and an *in vivo* unscheduled DNA synthesis (UDS) assay.

Tazobactam

Tazobactam was negative for mutagenic potential in an *in silico* screen. Tazobactam was negative for genotoxicity in an *in vitro* microbial mutagenicity (Ames) assay, an *in vitro* chromosomal aberration assay in Chinese hamster lung cells, a mammalian point-mutation (Chinese hamster ovary cell HPRT) assay, an *in vivo* rat chromosomal aberration assay, an *in vivo* mouse bone-marrow micronucleus assay, and a UDS assay. Tazobactam was positive for genotoxicity in an *in vitro* mouse lymphoma assay at ≥ 3000 mcg/mL.

Ceftolozane and Tazobactam

Ceftolozane and tazobactam (2:1 ratio) was negative for genotoxicity in an *in vitro* mouse lymphoma assay and an *in vivo* rat bone-marrow micronucleus assay. In an *in vitro* chromosomal aberration assay in Chinese hamster ovary cells, ceftolozane and tazobactam (2:1 ratio) was positive for structural aberrations, but only at highly toxic concentrations.

Carcinogenicity

Long-term carcinogenicity studies in animals have not been conducted with ceftolozane and tazobactam, ceftolozane, or tazobactam.

Reproductive Toxicity

Ceftolozane

Ceftolozane had no adverse effect on fertility in male or female rats following once daily IV administration at doses up to 1000 mg/kg/day; providing a safety margin of approximately 3-fold, based on plasma AUC.

Embryo-fetal development studies performed with intravenous, once daily ceftolozane in mice and rats with doses up to 2000 and 1000 mg/kg/day, respectively, revealed no evidence of harm

to the fetus. The mean maternal plasma exposure (AUC) values associated with these doses were approximately 7 (mouse) and 4 (rat) times the mean clinical ceftolozane exposure in adults at the clinical dose of 1 g thrice-daily.

Pre-postnatal studies of ceftolozane: In animal studies ceftolozane was associated with impaired auditory startle response in rat pups. See **WARNINGS AND PRECAUTIONS**, Pregnant Women.

Tazobactam

In a rat fertility and general reproduction study with intraperitoneal tazobactam twice-daily, concurrent with maternal toxicity, slight decreases in implantation and resultant slight decreases in the live litter size and increased stillbirths, and reversible delays in renal development were observed at 640 mg/kg/day (approximately 4 times the recommended clinical daily dose based on body surface comparison). Male and female fertility parameters, fertility of F1 generation and embryonic development of F2 generation were not impaired at doses less than or equal to 640 mg/kg/day.

In an embryo-fetal study in rats, tazobactam administered intravenously at doses up to 3000 mg/kg/day (approximately 19 times the recommended human dose based on body surface area comparison) produced maternal toxicity (decreased food consumption and body weight gain) but was not associated with fetal toxicity. In rats, tazobactam was shown to cross the placenta. Concentrations in the fetus were less than or equal to 10% of those found in maternal plasma.

Pre-postnatal studies of tazobactam: In animal studies tazobactam was associated with peri/postnatal effects in rat pups. See **WARNINGS AND PRECAUTIONS**, Pregnant Women.

Other Toxicity Studies

Ceftolozane was assessed for phototoxic, antigenic and immunotoxic potential. Clinically relevant findings were limited to hypersensitivity potential under sensitized conditions.

REFERENCES

1. Solomkin J, Hershberger E, Miller B. Ceftolozane/Tazobactam Plus Metronidazole for Complicated Intra-abdominal Infections in an Era of Multidrug Resistance: Results From a Randomized, Double-Blind, Phase 3 Trial (ASPECT-cIAI). *Clin Infect Dis*. 2015 May 15;60(10):1462-71. doi: 10.1093/cid/civ097. Epub 2015 Feb 10.
2. Wagenlehner FM, Umeh O, Steenbergen J, et. al. Ceftolozane-tazobactam compared with levofloxacin in the treatment of complicated urinary-tract infections, including pyelonephritis: a randomised, double-blind, phase 3 trial (ASPECT-cUTI). *Lancet*. 2015 May 16;385(9981):1949-56. doi: 10.1016/S0140-6736(14)62220-0. Epub 2015 Apr 27.
3. Clinical and Laboratory Standards Institute (CLSI). *Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria that Grow Aerobically; Approved Standard – Ninth Edition*. CLSI document M07-A9, Clinical and Laboratory Standards Institute, 950 West Valley Road, Suite 2500, Wayne, Pennsylvania 19087, USA, 2012.
4. Clinical and Laboratory Standards Institute (CLSI). *Performance Standards for Antimicrobial Disk Diffusion Susceptibility Tests; Approved Standard – Eleventh Edition*. CLSI document M02-A11, Clinical and Laboratory Standards Institute, 950 West Valley Road, Suite 2500, Wayne, Pennsylvania 19087, USA 2012.
5. Clinical and Laboratory Standards Institute (CLSI). *Methods for Antimicrobial Susceptibility Testing of Anaerobic Bacteria; Approved Standard – Eighth Edition*. CLSI document M11-A8. Clinical and Laboratory Standards Institute, 950 West Valley Road, Suite 2500, Wayne, Pennsylvania 19087, USA, 2012.
6. Clinical and Laboratories Standards Institute (CLSI). *Performance Standards for Antimicrobial Susceptibility Testing; Twenty-fourth Informational Supplement*. CLSI document M100-S24 [ISBN 1-56238-897-5 [Print; ISBN 1-56238-898-3 [Electronic]]. Clinical and Laboratory Standards Institute, 950 West Valley Road, Suite 2500, Wayne, Pennsylvania 19087-1898 USA, 2014.

**READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE
PATIENT MEDICATION INFORMATION**

 **ZERBAXA**[®]

zer bax' ah

ceftolozane and tazobactam

Read this carefully before you start taking **ZERBAXA**[®]. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **ZERBAXA**[®].

Serious Warnings and Precautions

Severe and sometimes fatal allergic reactions (e.g. anaphylaxis) have occurred with beta-lactam antibiotics. Tell your doctor if you are allergic to antibacterial drugs such as penicillins, cephalosporin or carbapenem. If you have a severe allergic reaction, stop taking ZERBAXA[®] and get immediate medical attention (See What are the possible side effects).

What is ZERBAXA[®] used for?

ZERBAXA[®] is used by healthcare professionals to treat complicated infections within the abdominal cavity and urinary tract infections, including a condition called “pyelonephritis” (a type of urinary tract infection that affects one or both kidneys) in adults.

Antibacterial drugs like ZERBAXA[®] treat only bacterial infections. They do not treat viral infections such as the common cold. Although you may feel better early in treatment, ZERBAXA[®] should be taken exactly as directed. Misuse or overuse of ZERBAXA[®] could lead to the growth of bacteria that will not be killed by ZERBAXA[®] (resistance). This means that ZERBAXA[®] may not work for you in the future. Do not share your medicine.

How does ZERBAXA[®] work?

ZERBAXA[®] contains 2 active substances. Ceftolozane prevents the growth of the bacterial cell wall and tazobactam binds to bacterial enzymes (e.g. beta-lactamases) that breakdown the antibiotics. Both work together to kill bacteria and reduce the infection.

What are the ingredients in ZERBAXA[®]?

Medicinal ingredients: ceftolozane 1 g (as ceftolozane sulfate) and tazobactam 0.5 g (as tazobactam sodium) per vial.

Non-medicinal ingredients: citric acid, L-arginine, and sodium chloride

ZERBAXA[®] comes in the following dosage forms:

ZERBAXA[®] is available as a lyophilized powder injection for intravenous use.

Do not use ZERBAXA[®] if you are hypersensitive (allergic) to:

- this drug or to any ingredient listed above.
- antibiotics like penicillin, or medicines known as “cephalosporins”, or other beta-lactams.

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take ZERBAXA[®]. Talk about any health conditions or problems you may have, including if you:

- know you are, or have previously been hypersensitive to penicillins, cephalosporins, beta-lactamases, or other antibacterial medicines.
- have recently had diarrhea, or have had diarrhea before taking this medicine.
- have liver or kidney problems.
- are pregnant or planning to become pregnant. If you think you may be pregnant or are planning to have a baby, ask your healthcare professional or pharmacist for advice before taking this medicine.
- are breast-feeding or planning to breastfeed.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

The following may interact with ZERBAXA[®]:

- Probenecid (a drug used to treat gout)

How to take ZERBAXA[®]:

The usual total daily dose of ZERBAXA[®] for adults is 1.5 g (1 g ceftolozane and 0.5 g tazobactam) administered every 8 hours by intravenous (IV) infusion over 1 hour, administered by the healthcare professional.

If you have kidney problems, your dose may be reduced.

Overdose:

If you think you have taken too much ZERBAXA[®], contact your healthcare professional, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.

Missed Dose:

ZERBAXA[®] is usually administered by a healthcare professional. If you suspect a missed dose, talk to your healthcare professional.

What are possible side effects from using ZERBAXA[®]?

Like all medicines, ZERBAXA[®] may have side effects. The most common side effects (may affect up to 1 in 10 people) are:

- Increase in the number of certain types of blood cells known as platelets
- Anemia
- Decrease in potassium (from blood tests)
- Insomnia
- Anxiety
- Headache

- Dizziness
- Atrial fibrillation (abnormal heart rhythm)
- Decrease in blood pressure
- Nausea
- Diarrhea
- Constipation
- Vomiting
- Abdominal pain (stomach ache)
- Rash
- Fever (high temperature)
- Increased liver enzymes (from blood tests)
- Local problems (e.g. abnormal redness of the skin, inflammation, pain, itching, or rash) when putting a substance into a vein (infusion site reactions)

Uncommon side effects (may affect up to 1 in 100 people) are:

- Inflammation of the large intestine due to *C. difficile* bacteria
- Inflammation of the stomach
- Abdominal bloating
- Indigestion
- Excessive gas in stomach or bowel
- Obstruction of the intestine
- Yeast infection in the mouth (thrush)
- Yeast infection of female genitalia
- Fungal urinary tract infection
- Increase in sugar (glucose) levels (from blood tests)
- Decrease in magnesium levels (from blood tests)
- Decrease in phosphate levels (from blood tests)
- Ischemic stroke (stroke caused by reduced blood flow in brain)
- Venous thrombosis (blood clot in a vein)
- Low red blood cell counts
- Atrial fibrillation (a condition involving rapid, irregular heartbeat)
- Fast heart beat
- Angina pectoris (chest pain or feeling of tightness, pressure or heaviness in chest)
- Itchy rash or swelling on the skin (hives)
- Kidney problems
- Kidney disease
- Shortness of breath

If you experience symptoms such as severe diarrhea (bloody or watery) with or without fever, abdominal pain, or tenderness, you may have Clostridium difficile colitis (bowel inflammation). If this occurs, stop taking ZERBAXA[®] and contact your healthcare professional immediately.

These are not all the possible side effects you may feel when taking ZERBAXA[®]. If you experience any side effects not listed here, contact your healthcare professional.

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
<u>RARE</u> Allergic Reaction: rash, hives (skin eruptions), swelling of the face, lips, tongue or throat, difficulty swallowing or breathing,			<u>✓</u>
Liver Disorder: yellowing of the skin or eyes, dark urine, pale stools		<u>✓</u>	

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, talk to your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<http://www.hc-sc.gc.ca/dhpmps/medeff/report-declaration/index-eng.php>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

Unopened vials: Store in a refrigerator (2°C – 8°C).

Store in the original package in order to protect from light.

Keep out of reach and sight of children.

If you want more information about ZERBAXA[®]:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the [Health Canada website](#) or Merck Canada website www.merck.ca or by calling Merck Canada at 1-800-567-2594.

To report an adverse event related to ZERBAXA[®], please contact 1-800-567-2594.

This leaflet was prepared by Merck Canada Inc.

Last Revised: July 26, 2017

® Merck Sharp & Dohme Corp. Used under license.

© 2017 Merck Canada Inc. All rights reserved.